### Level 3 Project Study Plan

### 2015 Big Creek Environmental Monitoring

(1) Objectives

In 2015, the Northeast Ohio Regional Sewer District (NEORSD) plans to conduct stream monitoring activities at six sites on Big Creek for the purpose of general watershed monitoring. The objective of the study is to monitor and assess the Big Creek watershed. The six sites will include River Miles (RM) 9.80, 4.40, and 0.15 on the Main Branch; RM 0.02 on the West Branch, RM 0.20 on an unnamed tributary that enters Big Creek Main Branch at RM 7.78; and Stickney Creek at RM 0.15.

RM 0.15 is located downstream of Jennings Road on the Big Creek Main Branch and is downstream of NEORSD-owned combined sewer overflows (CSOs). RM 4.40 is on Big Creek upstream of NEORSD CSOs and is in Memphis/Tiedeman Park. An assessment of stream habitat, water chemistry and fish and macroinvertebrate health at RM 0.15 and RM 4.40 are required by Ohio Environmental Protection Agency (Ohio EPA) National Pollutant Discharge Elimination System (NPDES) Permit No. 3PA0002\*GD.

RM 0.15 is also required to be monitored for water chemistry for the NEORSD consent decree monitoring plan under part of the Long Term Control Plan (LTCP) to assess stream improvement over time.

NEORSD will assess stream habitat, water chemistry, and fish and benthic macroinvertebrate community health to evaluate the impact of CSOs and other environmental factors on the creek at all six sites on Big Creek and Stickney Creek.

Fish and macroinvertebrate community health will be evaluated through the use of Ohio EPA's Index of Biotic Integrity (IBI), Modified Index of Well-Being (MIwb), and Invertebrate Community Index (ICI). An examination of the specific characteristics of the biological communities will be used in conjunction with water quality data, the NEORSD Macroinvertebrate Field Sheet and Qualitative Habitat Evaluation Index (QHEI) results in order to identify impacts to the communities. Results will be compared to historic data to show temporal as well as spatial trends. Water chemistry data will also be compared to the Ohio Water Quality Standards to determine attainment of applicable uses (Ohio EPA, 2011)<sup>1</sup>.

### (2) Point/Nonpoint Sources

<sup>&</sup>lt;sup>1</sup> See Appendix H for a list of references.

Point Sources	Nonpoint Sources
Combined Sewer Overflows	Urban Runoff
Storm Sewer Outfalls	Landfill leachate
Sanitary Sewer Overflows	Spills
Septic Tanks	Agriculture
NPDES Permitted Locations	

### (6) Sampling Locations

The following sample locations will be surveyed on Big Creek during the 2015 field season. Benthic macroinvertebrate and water chemistry collection sites are located near the midpoint of each electrofishing zone, indicated by RM. GPS coordinates are recorded at the downstream end of each electrofishing zone.

Water Body	Latitude	Longitude	River Mile	Location Information	USGS HUC 8 Number Name	Purpose
Big Creek, Main Branch	41.4460	-81.6865	0.15	Downstream of Jennings Road	04110002 Cuyahoga	Evaluate water chemistry, macroinvertebrates, fish and instream habitat as required by Ohio EPA Permit 3PA0002*GD
Big Creek, Main Branch	41.4460	-81.7540	4.40	Memphis Avenue Memphis Tiedeman Park	04110002 Cuyahoga	Evaluate water chemistry, macroinvertebrates, fish and instream habitat as required by Ohio EPA Permit 3PA0002*GD
Big Creek, West Branch	41.4459	-81.7545	0.02	Memphis Avenue Memphis Tiedeman Park	04110002 Cuyahoga	Evaluate water chemistry, macroinvertebrates, the fish community and instream habitat
Stickney Creek	41.4394	-81.7494	0.15	South of Memphis Ave.& north of Memphis Villas Blvd.	04110002 Cuyahoga	Evaluate water chemistry, macroinvertebrates, the fish community and instream habitat
Big Creek, Unnamed Tributary	41.4089	-81.7511	0.20	Upstream on Big Creek Parkway, Snow Rd. & Pearl Rd. Branch	04110002 Cuyahoga	Evaluate water chemistry, macroinvertebrates the fish community and instream habitat
Big Creek, Main Branch	41.3885	-81.7659	9.80	Downstream of Pearl Road	04110002 Cuyahoga	Evaluate water chemistry, macroinvertebrates, the fish community and instream habitat





### Big Creek Study Plan



### Legend

- Monitoring Site
- Regional Drainage
- CSO Outfall
- District Facility
- Outfalls
- --- NEORSD CSO Combined Sewer
- NEORSD CSO Responsibility Sewer
- NEORSD Intercommunity Relief Sewer
- NEORSD INTERCEPTOR
- Local Combined Sewer
- Local Culverted Stream
- Local Sanitary Sewer
- Local Storm Sewer



This information is for display purposes only. The Northeast Ohio Regional Sewer District (NEORSD) makes no warranties, expressed or implied, with respect to the accuracy of and the use of this map for any specific purpose. This map was created to serve as base information for use in Geographic Information Systems (GIS) for a variety of planning and analysis purposes. The NEORSD expressly disclaims any hability that may result from the use of this map. For more information, please contact: NEORSD GIS Services, 3900 Euclid Avenue, Cleveland, Ohio 44115 ----(216) 881-6600 ---GIS@neorsd.org

### 2015 Project Study Plans

### (3) Parameters Covered

Fish specimens will be identified to species level, weighed, counted and examined for the presence of external anomalies including DELTs (deformities, eroded fins, lesions and tumors). An Ohio EPA Fish Data Sheet (Appendix A) will be completed during each assessment. Quantitative fish sampling is expected to be conducted at all locations.

Macroinvertebrate community assemblages will be collected from each location. Third Rock Consultants, LLC will identify and enumerate the specimens collected from each site. All specimens will be identified to the lowest practical taxonomic level as recommended in Ohio EPA's *Biological Criteria for the Protection of Aquatic Life, Volume III* (1987b)<sup>1</sup>. The NEORSD Macroinvertebrate Field Sheet (Appendix A) will be completed at each site during sampler retrieval or when qualitative sampling is conducted.

Stream habitat will be measured by scoring components of the QHEI at all locations, including the substrate, instream cover, channel morphology, riparian zone, bank erosion, pool/glide and riffle/run quality and gradient. The HHEI will be conducted at those sites with drainage areas less than one square mile listed under PSPs with general watershed monitoring. The Lacustuary QHEI (L-QHEI) will be performed at sites that are affected by the water level of Lake Erie. Examples of the Ohio EPA field sheets for the QHEI, L-QHEI and the HHEI can be found in Appendix A.

Water chemistry samples will be collected at each electrofishing/ macroinvertebrate sampling site included in the study. Water chemistry samples will be analyzed by NEORSD's Analytical Services Division. Appendix B lists the parameters to be tested along with the detection limits and practical quantitation limits. Field measurements for dissolved oxygen, pH, temperature, conductivity and turbidity will also be performed. A Surface Water Condition Sampling Field Data Form will be completed at each site during each sampling event (Appendix A).

Benthic and water column chlorophyll *a* samples may be collected from stream locations. Chemical and physical water quality parameters to be measured in conjunction with the chlorophyll *a* samples include total phosphorus, dissolved reactive phosphorus, nitrite, nitrate+nitrite, ammonia, alkalinity, turbidity and suspended solids. In the Cuyahoga River, YSI 6600EDS data sondes may be

<sup>&</sup>lt;sup>1</sup>See Appendix H for a list of all references.

installed at RMs 16.20, 10.75, 10.10, and 7.00 around the time that this sampling is conducted to more frequently monitor dissolved oxygen, temperature, conductivity, specific conductivity and pH.

### (4) Field Collection and Data Assessment Techniques

Field collections for fish will be conducted at all stream locations. Sampling will be conducted using longline, backpack, or boat electrofishing techniques and will consist of shocking all habitat types within a sampling zone. Headwater and wading sites, which are 0.15 and 0.20 kilometers in length, respectively, will be surveyed by moving from downstream to upstream. Boat sites, which are 0.50 kilometers in length, will be surveyed by moving from upstream to downstream. The stunned fish will be collected and placed into a live well for later identification. The longline, backpack, and boat electrofishing zones will be assessed one to three times during the field season (June 15 - October 15).

Fish will be identified to the species level, weighed, counted, and examined for the presence of external anomalies including DELTs. Fish easily identified (commonly collected from year to year) will be returned to the site from which they are collected. Fish species difficult to identify will be brought back to the laboratory for verification by NEORSD Level 3 Fish Qualified Data Collectors (QDC). If necessary, vouchers will be sent to The Ohio State University Museum of Biological Diversity for verification by the Curator and/or Associate Curator of Fish. Voucher specimens will be collected as described in section (14). Endangered species and those too large for preservation will not be collected as voucher specimens, but will instead be photographed. Photographed vouchers will include features that permit definitive identification of the particular species.

Fish will be preserved in 10 percent formalin in the field, soaked in tap water for 24 to 48 hours after 5 to 7 days, then transferred to solutions of 30 and 50 percent ethanol for 5 to 7 days each and, finally, to 70 percent ethanol for long-term storage. Specimens larger than six inches will be slit along the right side and then soaked in formalin for approximately 10 to 14 days before being transferred to water and solutions of 30, 50 and 70 percent ethanol. Label information will include location (description and coordinates), date, time, collectors' names and sample identification code for each specimen collected.

Macroinvertebrate sampling will be conducted using quantitative and qualitative sampling techniques. Quantitative sampling will be done using a modified Hester-Dendy multi-plate artificial substrate sampler (HD) that is colonized for a six-week period. Multiple HD samplers may be installed at one or all sampling locations in case samplers are lost due to vandalism, burial, etc. and for the purposes of providing a replicate sample. Qualitative sampling will be conducted using a D-frame dip net when HD samplers are retrieved. The NEORSD

Macroinvertebrate Field Sheet will be completed during each HD retrieval. Ronald Maichle of NEORSD, a Level 3 QDC for Benthic Macroinvertebrate Biology, may identify specimens in the replicate sample to the lowest practical taxonomic level as recommended in Ohio EPA's *Biological Criteria for the Protection of Aquatic Life, Volume III* (1987b).

Macroinvertebrate voucher specimens for both quantitative and qualitative sampling will be collected as described in section (14). Macroinvertebrate community assemblages collected will be shipped to Third Rock Consultants, LLC (Lexington, KY) for identification and enumeration. Third Rock Consultants, LLC will identify specimens to the lowest practical taxonomic level as recommended in Ohio EPA's *Biological Criteria for the Protection of Aquatic Life, Volume III* (1987b).

A detailed description of the sampling and analysis methods utilized in the fish community and macroinvertebrate surveys, including calculations of the IBI, MIwb, and ICI, can be found in Ohio EPA's *Biological Criteria for the Protection of Aquatic Life, Volumes II* (1987a) and *III* (1987b). Methods for assessing fish and macroinvertebrate communities in lacustuary zones can be found in Ohio EPA's draft *Biological Criteria for the Protection of Aquatic Life, Volume IV* (1997).

The QHEI, as described in Ohio EPA's *Methods for Assessing Habitat in Flowing Waters: Using the Qualitative Habitat Evaluation Index (QHEI)* (2006) will be used to assess aquatic habitat conditions at each sample location. The L-QHEI will be used where appropriate and will follow Ohio EPA's draft *Methods of Assessing Habitat in Lake Erie Shoreline Waters Using the Qualitative Habitat Evaluation Index (QHEI) Approach (Version 2.1)* (2010).

The HHEI as described in Ohio EPA's *Field Evaluation Manual for Ohio's Primary Headwater Habitat Streams* (2012a) will be used, when necessary, to conduct use attainability analyses and to classify the actual and expected biological conditions in primary headwater habitat streams.

Water chemistry sampling may occur across a variety of flow conditions. Techniques used for water chemistry sampling and chemical analyses will follow the *Surface Water Field Sampling Manual* (Ohio EPA, 2013). Chemical water quality samples from each site will be collected with at least one 4-liter disposable polyethylene cubitainers with disposable polypropylene lids and two 473-mL plastic bottles. Water samples collected for analysis of dissolved reactive phosphorus will be filtered using a 0.45-µm PVDF syringe filter and will be collected in a 125-mL plastic bottle. Bacteriological samples will be collected in a sterile plastic bottle preserved with sodium thiosulfate. All water quality samples

will be collected as grab samples. Field blanks and duplicate samples will each comprise not less than 5% of the total samples collected for this study plan, for a total frequency of quality control samples of not less than 10% of the total samples collected. With the exception of bacteriological duplicate samples, the acceptable percent RPD will be based on the ratio of the sample concentration and detection limit (Ohio EPA, 2013): Acceptable % RPD =  $[(0.9465X^{-0.344})*100] + 5$ , where X = sample/detection limit ratio. For bacteriological duplicates, duplicate samples more than 5x apart from one another ( $\[MPD > 133.3\]$ ) will be rejected in accordance with the Ohio EPA approved method for data validation of bacteriological samples outlined in Section F of the Ohio 2012 Integrated Water Quality Monitoring and Assessment Report (Ohio EPA, 2012b). Those RPDs that were higher than acceptable may indicate potential problems with sample collection and, as a result, the data will not be used for comparison to the water quality standards. Acid preservation of the samples, as specified in the NEORSD laboratory's standard operating procedure for each parameter, will occur in the field. Appendix B lists the analytical method, method detection limit and practical quantitation limit for each parameter analyzed. Field analyses include the use of either a YSI-556 MPS Multi-Parameter Water Quality Meter, YSI EXO1 sonde, or YSI 600XL sonde to measure dissolved oxygen (DO), water temperature, conductivity and pH; and when necessary, a Hanna HI 98129 meter to measure pH and a Hach HQ30d meter with LDO101 probe to measure DO. Field turbidity will be measured using either a Hach 2100P Portable Turbidimeter or Hach 2100Q Turbidimeter. Specifications for these meters have been included in Appendix C.

Benthic and water column chlorophyll a samples may be collected if time and resources allow. Sampling methods will follow those detailed in the NEORSD Chlorophyll a Sampling and Field Filtering Standard Operating Procedure (SOP-EA001-00). A Chlorophyll a Sampling Field Sheet will be completed for each site (Appendix D). Water chemistry grab samples will be collected at the same time using the methods discussed previously and will be analyzed for nutrients, turbidity, alkalinity and suspended solids. Additionally in the Cuyahoga River, approximately 24-hours prior to each chlorophyll a sampling event, YSI 6600 EDS data sondes may be deployed at RMs 16.20, 10.75, 10.10 and 7.00. If installed, each data sonde will record, at fifteen-minute intervals, dissolved oxygen concentration, pH, temperature, and conductivity from the time the data sonde is deployed until the time it is retrieved. These data sondes will be placed in the stream by inserting each one into a 4.5-inch PVC pipe with holes drilled into the sides of the lower third of the pipe to allow water to pass through it. The data sondes will remain in the river for approximately 24-hours or longer following collection of the chlorophyll *a* samples.

Where possible, data assessment will include an analysis of temporal and spatial trends in the collected data. Species assemblages and individual metrics will be

analyzed. Graphs that show current and historic QHEI, L-QHEI, IBI, LIBI, MIwb, ICI, and LICI scores and how these scores compare to attainment status of biocriteria will be prepared. Water chemistry data collected will be compared to Ohio water quality standards to determine whether any excursions from the applicable water quality criteria have occurred. It will also be used to determine any relationships among individual parameters and chlorophyll *a* concentrations. Comparisons between water quality and biological community health will only be made if at least three water quality samples have been collected from that site.

(5) Stream Flow Measurement

Stream flow will be recorded for all locations during each electrofishing pass utilizing data from the United States Geological Survey (USGS) gauge station nearest the stream location, if applicable.

Stream flow will be measured with a Marsh-McBirney FloMate Model 2000 Portable Flow Meter, a HACH FH950 Flow Meter or an Aquaflow Probe Model 6900, which measure flow in feet per second, when HD samplers are installed and retrieved. The specifications for the flow meters can be found in Appendix C.

(7) Schedule

One to three electrofishing surveys will be conducted at each site between June 15 and October 15, 2015. Surveys will be conducted at least three weeks apart. Specific dates have not been scheduled. River flow and weather conditions will be assessed weekly to determine when each electrofishing pass will be conducted.

Artificial substrate samplers will be installed at stream locations between June 15 and August 19, 2015, and retrieved six weeks later. Qualitative macroinvertebrate sampling will be conducted one time at all sites. Specific dates have not been scheduled. River flow and weather conditions will be assessed weekly to determine when the HD sampler installations and retrievals and qualitative sampling will be conducted.

QHEI, and if necessary, HHEI and L-QHEI habitat evaluations will be conducted one time between June 15 and October 15, 2015. QHEI evaluations will be conducted around the same time as one of the electrofishing surveys.

Water chemistry samples will be collected a minimum of three times from stream locations between June 15 and October 15, 2015.

Benthic and water column chlorophyll *a* samples may be collected at least one time from stream locations between June 15 and October 15, 2015. These samples will be collected under low-flow conditions.

### (8) QA/QC

Quality assurance and quality control of sampling and analysis methods for habitat, fish, and macroinvertebrate evaluations will follow Ohio EPA's *Biological Criteria for the Protection of Aquatic Life, Volumes II* (1987a) and *III* (1987b), *Methods for Assessing Habitat in Flowing Waters: Using the Qualitative Habitat Evaluation Index (QHEI)* (2006), *Field Evaluation Manual for Ohio's Primary Headwater Habitat Streams* (2012a), draft *Biological Criteria for the Protection of Aquatic Life: Volume IV: Fish and Macroinvertebrate Indicies for Ohio's Lake Erie Nearshore Waters, Harbors, and Lacustuaries* (1997) and draft *Methods of Assessing Habitat in Lake Erie Shoreline Waters Using the Qualitative Habitat Evaluation Index (QHEI) Approach (Version 2.1)* (2010)

Electrofishing equipment will be used according to the guidelines listed in the operation and maintenance manual provided by Smith-Root, Inc. Malfunctioning equipment will not be used to collect data. Proper steps will be taken to correct any problems as soon as possible, whether by repairing in the field, at the NEORSD Environmental & Maintenance Services Center, or by contacting the supplier or an appropriate service company.

Fish species difficult to identify will be brought back to the laboratory for verification by Level 3 Fish QDC's, and if necessary, sent to The Ohio State University Museum of Biological Diversity for verification by the Curator and/or Associate Curator of Fish. Voucher specimens will be collected as described in section (14). Endangered species and those too large for preservation will not be collected as voucher specimens, but will instead be photographed. Photographed vouchers will include features that permit definitive identification of the particular species.

All macroinvertebrate community assemblages from stream locations, except for the replicate sample, will be collected and shipped to Third Rock Consultants, LLC for identification and enumeration. All specimens will be identified to the lowest practical taxonomic level as recommended in Ohio EPA's *Biological Criteria for the Protection of Aquatic Life, Volume III* (1987b). All macroinvertebrate specimens will be returned to NEORSD. At least two voucher specimens of each species, when available, will be separated into individual vials and kept as described in section (14). The remaining specimens for each site will be returned in a single container labeled with the site number and collection method and date. All specimens and accompanying chain-of-custody documentation will be retained by NEORSD and stored at the Environmental & Maintenance Services Center for a period not less than ten years. Water samples obtained for chemical analyses will be collected, preserved (see Section 4), labeled and then placed on ice inside the field truck. The field truck will remain locked at all times when not occupied/visible. Sampling activities, including sample time and condition of surface water sampled, will be entered in a field log book and on the Surface Water Condition Sampling Field Data Form. The samples will then be delivered immediately to the NEORSD Analytical Services cooler, after which the door to the cooler will be locked, and the samples will be transferred to the custody of Analytical Services. The NEORSD Analytical Services Quality Manual and associated Standard Operating Procedures are on file with Ohio EPA. The Quality Assurance Officer at Analytical Services will send updates, revisions and any information on document control to Ohio EPA as needed.

For benthic and water column chlorophyll *a* sampling, three filtrations will be performed for each sample. A field filtration blank will be submitted for every 20 samples.

Calibration of YSI 6600EDS data sondes will be done according to the YSI Environmental Operations Manual. The conductivity will be calibrated first using a 1.413 mS/cm standard. Second, the pH will be calibrated using two different buffers (7 and 10 s.u.). The DO will be calibrated last with an acceptable error of 0.2 mg/L.

Once the sondes are removed from the river, the accuracy of the data that has been collected will be checked by comparing readings taken by the sondes to known standards. If the measurements taken at this time meet quality control goals, all of the data collected since the last calibration will be considered accurate. The acceptable differences for pH and conductivity will be  $\pm 0.3$  with pH 7 buffer and  $\pm 10\%$  of the conductivity standard, respectively (EPA New England- Region 1, 2005). The acceptable difference for DO will be  $\pm 0.2$  mg/L. If the measurements do not meet quality control goals, best professional judgment will be used to decide if any of the data collected during that period may still be accurate. For example, the data collected from the four locations may be plotted on the same graph, and if it appears that the data points are following similar trends, they may be considered accurate. If any data that do not meet quality control goals are used, a rationale for their inclusion will be provided when the data are submitted.

### (9) Work Products

Within one year of completion of the project, fish data (species, numbers, weights, pollution tolerances, the incidence of DELT anomalies, IBI or LIBI, MIwb scores), macroinvertebrate data (types and numbers of macroinvertebrates collected and ICI or LICI scores), habitat data (QHEI or L-QHEI raw data and scores) and water chemistry results will be submitted to the Ohio EPA or an Ohio

EPA approved data warehouse. Additionally, reports summarizing, interpreting, graphically presenting and discussing the IBI (LIBI, where applicable), MIwb, ICI (LICI, where applicable) and QHEI (L-QHEI, where applicable) scores, chlorophyll *a* results, and any excursions from water quality standards may be prepared for internal use.

### (10) Qualified Data Collectors

The following Level 3 Qualified Data Collectors (QDC) will be involved with this study:

Name	Address	Email Address	Phone Number	QDC Specialty(s)
John W. Rhoades <sup>1</sup>	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	rhoadesj@neorsd.org	216-641-6000	QDC - 00008 CWQA/FCB/SHA/ BMB
Cathy Zamborsky	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	zamborskyc@neorsd.org	216-641-6000	QDC - 00009 CWQA/SHA
Seth Hothem	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	hothems@neorsd.org	216-641-6000	QDC - 00010 CWQA/FCB/SHA/ BMB
Tom Zablotny	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	zablotnyt@neorsd.org	216-641-6000	QDC - 00018 CWQA/FCB/SHA
Ron Maichle	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	maichler@neorsd.org	216-641-6000	QDC - 00145 CWQA/SHA/BMB
Francisco Rivera	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	riveraf@neorsd.org	216-641-6000	QDC - 00262 CWQA/SHA
Jillian Knittle	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	knittlej@neorsd.org	216-641-6000	QDC – 00512 CWQA/SHA/BMB
Jonathan Brauer <sup>2</sup>	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	brauerj@neorsd.org	216-641-6000	QDC - 00663 SHA
Bert Remley <sup>3</sup>	2526 Regency Road, Suite 180 Lexington, Kentucky 40503	bremley@thirdrockconsultants.com	859-977-2000	QDC - 00837 BMB
NEORSD Lead Proje	ect Manager			
<sup>2</sup> See acknowledgemer	at letter for conducting QHEIs (App	endix F)		
Benthic Macroinverte	brate Identification			

The following is a list of persons not qualified as Level 3 QDCs who may be involved in the project. Prior to the start of sampling, the project managers will explain to each individual the proper methods for sampling. Sampling will only be completed under the direct observation of a QDC. The lead project manager will be responsible for reviewing all reports and data analysis prepared by qualified personnel prior to completion.

Name	Address	Email Address	Phone Number
Kelsey Amidon	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	amidonk@neorsd.org	216-641-6000
Nick Barille	4747 East 49 <sup>th</sup> Street	barillen@neorsd.org	216-641-6000

### PSP Guidelines 3-5 & 7-17 April 6, 2015

Name	Address	Email Address	Phone Number
	Cuyahoga Hts., Ohio 44125		
Mark Colvin	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	colvinm@neorsd.org	216-641-6000
Tim Dobriansky	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	dobrianskyt@neorsd.org	216-641-6000
Donna Friedman	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	friedmand@neorsd.org	216-641-6000
Rae Grant	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	grantr@neorsd.org	216-641-6000
Mark Matteson	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	mattesonm@neorsd.org	216-641-6000
Mario Meany	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	meanym@neorsd.org	216-641-6000
Carrie Millward	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	millwardc@neorsd.org	216-641-6000
Denise Phillips	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	phillipsd@neorsd.org	216-641-6000
Brandy Reischman	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	reischmanb@neorsd.org	216-641-6000
Frank Schuschu	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	schuschuf@neorsd.org	216-641-6000
Eric Soehnlen	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	soehnlene@neorsd.org	216-641-6000
William Stanford	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	standfordw@neorsd.org	216-641-6000
Wolfram von Kiparski	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	vonkiparskiw@neorsd.org	216-641-6000
NEORSD Summer Co-op #1	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	To Be Determined	216-641-6000
NEORSD Summer Co-op #2	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	To Be Determined	216-641-6000
NEORSD Summer Co-op #3	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	To Be Determined	216-641-6000
NEORSD Summer Co-op #4	4747 East 49 <sup>th</sup> Street Cuyahoga Hts., Ohio 44125	To Be Determined	216-641-6000

### (11) Contract laboratory contact information

All bacteriological and/or chemical sample analysis will be completed by NEORSD's Analytical Services Division. Evidence of NEORSD's Analytical Services current accreditation and method dates can be found in Appendix E. The contact information for NEORSD's Analytical Service Division is:

NEORSD Analytical Services Mr. Mark Citriglia 4747 E. 49th Street Cuyahoga Heights, Ohio 44056 citrigliam@neorsd.org 216-641-6000 Any fish that is not positively identified in the field, or at NEORSD, will be sent to The Ohio State University Museum of Biological Diversity for verification by the Curator and/or Associate Curator of Fish. Fish will be identified to the species level.

Dr. Ted Cavender, Curator of Fish / Mr. Marc Kibbey, Associate Curator of Fish 1315 Kinnear Road, Columbus, Ohio 43212 cavender.1@osu.edu / kibbey.3@osu.edu 614-292-7873

Identification of macroinvertebrates for stream locations will be completed by Third Rock Consultants LLC (Lexington, Kentucky) (Appendix F). Benthic macroinvertebrates will be identified to the lowest practical level as recommended by Ohio EPA (1987b). Third Rock Consultants LLC contact information:

Ms. Marcia Wooton Third Rock Consultants LLC 2526 Regency Road, Suite 180 Lexington, Kentucky 40503 <u>mwooton@thirdrockconsultants.com</u> 859-977-2000

- (12) Copy of ODNR collector's permit See Appendix G.
- (13) Digital Catalog Statement

A digital photo catalog of all sampling locations will be maintained for 10 years and will include photos of the specific sampling location(s), the riparian zone adjacent to the sampling location(s) and the general land use in the immediate vicinity of the sampling location(s).

- Date: 04/13/15 Ton Print/Signature: John W. Rhoades

(14) Voucher Specimen Statement

NEORSD will maintain a benthic macroinvertebrate and fish voucher collection which includes two specimens, or appropriate photo vouchers, of each species or taxa collected during the course of biological sampling from any stream within the NEORSD's service area. When benthic macroinvertebrates from multiple surface waters are collected within the same year and identified by the same QDC, one voucher collection will be created to represent the specimens collected from those streams. When fish specimens from multiple surface waters are collected within the same year, one voucher collection will be created to represent the specimens collected from those streams. A separate collection for each sampling event will not be maintained.

NEORSD will provide specimens or photo vouchers to the Director upon request. This collection will be stored at the NEORSD laboratory in the Environmental and Maintenance Services Center.

Print/Signature: John W. Rhoades

Date: 04/13,

(15) Sample Location Statement

I attest that I will make available any and all sampling location information, including but not limited to; the name of the water body sampled, sampling location latitude and longitude, sampling location river mile where possible, general location information, the U.S. geological survey HUC 8 number and name, and the purpose for data collection at each sampling location.

Print/Signature: John W. Rhoades 4 Date: 04

(16) Additional L3 Data Collector Statement

The Lead Project Manager for all stream locations is approved for all project data types.

Date: 04 Print/Signature: John W. Rhoades

### (17) Trespassing Statement

I have not been convicted or pleaded guilty to a Violation of section 2911.21 of the Revised Code (criminal trespass) or a substantially similar municipal ordinance within the previous five years

	alt		, 1
Print/Signature:	John W. Rhoades	Date:	04/13/15
Print/Signature:	Cathy Zamborsky / Tthy Josnoly	Date:	4/10/15
Print/Signature:	Seth Hothem / Ber Notes	Date:	4/10/15
Print/Signature:	Tom Zablotny / Jon Zary	Date:	41015
Print/Signature:	Ron Maichle Mark Munitor	Date:	04-10-15
Print/Signature:	Jillian Knittle / Jeliffelter	Date:	4/13/15
Print/Signature:	Francisco Rivera / Francin ) This	Date:	4/13/15
Print/Signature:	Jonathan Brauer / Mar	Date:	4/10/15
	10		

Appendix A

Chiezza	SHEET		(1	requires lat/long & cou	nty)	Lone		Pag	ge	of	
Station ID		_ River Code_		RM	Date			_Tiı	ne	_	
Stream				Location	I						
Comments —				<u> </u>							
Lat	Lon	ng	County _		ALP _		_ Tir	ne F	ishe	d	
Crew		_Netter	Othe	rs		Sam	pler	Туре			
Distance	Flow	Temp. C	Secchi	Source	Project_						
Fins Code	Number To Weighed Co	otal Total unted Weight		Weights	ounts	Defor	DE mities Multi	LT A , Eros ple DI	NOM ions, I ELTs c	IALI Lesion	ES 1s, Tum fish
						D	E	L	Т	М	*
		· · · · ·									
V 10v						-					<u> </u>
						D	E	L	Т	M	*
V 10-											
• IUX						D	E	L	Т	M	*
	- <b> </b>										
						_					
V 10x							E	T	т	M	*
							E		1	141	
V 10x							17				
						D	E	L 	1.	M	<b></b>
V 10x											
						D	E	L	Т	М	ale
V 10x											+
						D	E	L	Т	М	*
	1										
V 10-	,					- _			ļ	<u> </u>	-
• 10x						D	E	L	Т	М	*
	l	I					-				+
V 10x		1				D	E	L	Т	М	*
							-	[ <sup>-</sup>			-
V 10x		-						1	1	1	1

\* A-anchor worm; B-black spot; C-leeches; F-fungus; N-blind; P-parasites; S-emaciated, W-swirled scales Y-popeye; Z-other

11/4/2005



Stream				Dim	» Mile:		V	
Stream:				Kive			r ear:	
Location:	2		¥	Project:				
Drainage Area (n	ni²):	Latitud	te (°N)/Longitud	le (°W):				
			Hester-Dendy	Deploymen	t Informatic	n		
Install Date:			-	Crew (QDC (	Circled):			
Current at HD (f	ps):		Depth (cn	ı):		Pictures	Obtained: Yes	No
Reinstall Date:				Crew (QDC (	Circled):			
Current (fps):		Depth (	cm):		Reason:			
Reinstall Date:				Crew (QDC (	Circled):			
Current (fps):		Depth (	cm):		Reason:			
			Sampling/	Retrieval In	formation			
Sampling Methor	d:	Hester-Dend	ly Dipn	et Surb	er Coi	re Oth	ner:	
Sample ID	: HD	:	Qua	litative:		Other		
Sampling Date:			Crew	(QDC Circle	ed):			
HD Condition-	Current	(fns):	Dont	) (cm):		Water Temp		°F / °C
The Condition-	Number	r of HD Block	s Obtained	. (em).	Rem	arks:	·	170
	Disturb	ed: Ye	s No (	Comments:				
	Debris:	Ye	s No (	Comments:				
	Silt/Sol	ids: No	ne Sligh	t Mod	erate	Heavy		
Dipnet-	Time Sa	ampled (min):		X Number of	of Crew:	= To	tal (min):	
-	Habitat	s Sampled:	Pool I	Riffle	Run	Margin	Backwater	
			River S	ampling Cor	ditions			
Flow Condition:		Flood	Above Normal	Normal	Low	Interstitial	Intermittent	Dry
Current Velocity.	:	Fast	Moderate	Slow	Non-det	ect		
Channel Morpho	logy:	Natural	Channelized	Channeli	zed (Recover	red) Imp	oounded	
Bank Erosion:		Extensive	Moderate	Slight	None			
Riffle Developme	ent:	Extensive	Moderate	Sparse	Absent			
Riffle Quality:		Good	Fair	Poor		Embedded:	Yes	No
Water Clarity:		Clear	Murky	Turbid	-	Other:		
Water Color:		None	Green	Brown	Grey	Other:		
Canopy over HD	•	Open	13 70	JU %	23 %	Closed		
Comment Section	on:							
OEPA Commen	t Field C	Codes:						

Last Modified 01/31/13

Substrate C	Characte	eristic	s		•	Predominant La	nd Use (Lef	it, Right or Both	ı)
	-	e		ſ		Forest	Urban		Open Pasture
	Pool	u Hi	its	Rur	its	Shrub	Residential	/Park	Closed Pasture
		- <b>H</b>	CP		Un	Old Field	Mining/Con	nstruction	
Bedrock						Rowcrop	Wetland		
Boulder						Industrial	Other		
Rubble									
Coarse Gravel						Predominant Ri	parian Vege	etation	
Fine Gravel						Left	Right	Туре	
Sand								Large Tre	ees
Silt						Small Trees			ees
Clay/Hardpan								Shrubs	
Detritus								Grass/We	eds
Peat			-					None	
Muck		-				<u></u>			
Other		-				Margin Habitat			
Macrophytes		-				Margin Ouality:	Good	Fair	Poor
Algae						Undercut Ba	nks Ro	oot Mats	Tree Roots
Artifacts						Grass	W	ater Willow	Woody Debris
Compaction (F.M.S)			-			Shallows	C	av/Hardnan	Macrophytes
Denth (Avg)						Rin Ran	B	ulkhead	macrophytes
Width (Avg)	$\square$		-	$\vdash$		Other		anticad	
						<u> </u>			
					Biolo	gical Characteris	tics		
					DIVIO	gical Characteris	1163		
Riffle:					DIOIO	gical Characteris	V≃ Very	Abundant; A= Abunda	nt; C= Common; R= Rare
Riffle: Predominant Org	ganism:				Diolo	gital Characteris	V≃ Very Overall An	Abundant; A= Abundan 10unt (V=>1	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1)
Riffle: Predominant Org Other Common (	ganism: Organisr	ns:				gicai Characteris	V≃ Very Overall An	Abundant; A= Abundan nount (V=>1 Porifera, Bryozy	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) Da
Riffle: Predominant Org Other Common ( Density:	ganism: Organisr High	ns:N	лоder	ate	Low		V≃ Very Overall An / / /	Abundant; A= Abundan nount (V=>19 Porifera, Bryozo Turbellaria, Oli	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea
Riffle: Predominant Org Other Common ( Density: Diversity:	ganism: Organisr High High	ns:N N	/loder	ate	Low		V≃ Very Overall An / / /	Abundant; A= Abundan 10unt (V=>1 Porifera, Bryoz Turbellaria, Oli Isopoda, Amph	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda
Riffle: Predominant Org Other Common ( Density: Diversity:	ganism: Organisr High High	ns <u>:</u> N N	/loder /loder	ate ate	Low	,	V≃ Very Overall An / / / / /	Abundant; A= Abundan 10unt (V=>12 Porifera, Bryoz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) 0a gochaeta, Hirudinea ipoda racarina
Riffle: Predominant Org Other Common ( Density: Diversity: Run:	ganism: Organisr High High	ns:N N	/loder /loder	ate ate	Low		V≃ Very Overall An / / / /	Abundant; A= Abundan nount (V=>12 Porifera, Bryozo Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org	ganism: Organisr High High ganism:	ns:N N	/loder	ate ate	Low		V≃ Very Overall An / / / / /	Abundant; A= Abundan nount (V=>1 Porifera, Bryozz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common (	ganism: Organisr High High ganism: Organisr	ns:N N 	Лоder Лoder	ate ate	Low Low		V≃ Very Overall An / / / / / / / /	Abundant; A= Abundan nount (V=>15 Porifera, Bryozz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniid	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density:	ganism: Organisr High High ganism: Organisr High	ns:N N N	/loder /loder	rate rate	Low Low	, ,	V≃ Very Overall An / / / / / / /	Abundant; A= Abundan 10unt (V=>1) Porifera, Bryoz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity:	ganism: Organisr High High ganism: Organisr High High	ns:N N N N	Aoder Aoder Aoder	rate rate rate rate rate	Low Low Low Low	, ,	V≃ Very Overall An / / / / / /	Abundant; A= Abundan 10unt (V=>1) Porifera, Bryoz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity:	ganism: Organisr High High ganism: Organisr High High	ns:N N N N	Aoder Aoder Aoder Aoder	ate ate ate ate ate	Low Low Low Low	, ,	V≃ Very Overall An / / / / / / /	Abundant; A= Abundan nount (V=>12 Porifera, Bryoza Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniid Other Zygoptera, Anis Plecoptera	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool:	ganism: Organisr High High ganism: Organisr High High	ns:N N N N	Aoder Aoder Aoder Aoder	ate ate ate ate	Low Low Low Low	, ,	V≃ Very Overall An / / / / / / /	Abundant; A= Abundan nount (V=>12 Porifera, Bryoza Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org	ganism: Organisr High High ganism: Organisr High High ganism:	ns:N N N	Aoder Aoder Aoder Aoder	rate ate rate rate	Low Low Low Low	, ,	V≃ Very Overall An / / / / / / / / /	Abundant; A= Abundan nount (V=>12 Porifera, Bryozz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae  soptera
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common (	ganism: Organisr High High ganism: Organisr High High ganism: Organisr	ns:N N N N	Aoder Aoder Aoder Aoder	rate ate rate rate	Low Low Low Low	, ,	V≃ Very Overall An / / / / / / / / /	Abundant; A= Abundan nount (V=>1) Porifera, Bryozy Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae soptera
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density:	ganism: Organisr High High ganism: Organisr High High ganism: Organisr High	ns:N N N N	Aoder Aoder Aoder Aoder	rate ate rate rate	Low Low Low Low Low	, ,	V≃ Very Overall An / / / / / / / /	Abundant; A= Abundan nount (V=>15 Porifera, Bryozz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae soptera europtera
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Diversity:	ganism: Organisr High High ganism: Organisr High ganism: Organisr High High	ns:N N N N	Aoder Aoder Aoder Aoder Aoder	rate rate rate rate rate rate	Low Low Low Low Low Low	, , ,	V≃ Very Overall An / / / / / / / / / / /	Abundant; A= Abundan nount (V=>12 Porifera, Bryoza Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera Hydropsyc Hydropsyc	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae 
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Diversity:	ganism: Organisr High High ganism: Organisr High ganism: Organisr High High	ns:N N N 	Aoder Aoder Aoder Aoder Aoder	rate rate rate rate rate rate	Low Low Low Low Low Low	, , ,	V= Very Overall An / / / / / / / / / /	Abundant; A= Abundan 10unt (V=>12 Porifera, Bryoza Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera Hydropsyc Hydropsyc Other	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Diversity: Margin:	ganism: Organisr High High ganism: Organisr High ganism: Organisr High High	ns:N N N N N	Aoder Aoder Aoder Aoder Aoder	rate rate rate rate rate rate	Low Low Low Low Low Low	, , ,	V= Very Overall An / / / / / / / / / / / / /	Abundant; A= Abundan tount (V=>12 Porifera, Bryoze Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera Hydropsyc Hydropsyc Coleoptera	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Diversity: Margin: Predominant Org	ganism: Organisr High High ganism: Organisr High High High High High	ns:N N N N N	Aoder Aoder Aoder Aoder Aoder	ate rate rate rate rate rate	Low Low Low Low Low Low	, , ,	V≃ Very Overall An / / / / / / / / / / / / /	Abundant; A= Abundan tount (V=>12 Porifera, Bryozz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptagenito Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera Hydropsyc Hydropsyc Coleoptera Elimidae	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Margin: Predominant Org Other Common (	ganism: Organisr High High ganism: Organisr High High High High ganism: Organisr	ns:N Ns:N ns:N ns:N	Aoder Aoder Aoder Aoder Aoder	rate ate rate rate rate	Low Low Low Low Low Low	, , ,	V≃ Very Overall An / / / / / / / / / / / / /	Abundant; A= Abundan nount (V=>12 Porifera, Bryozz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera Hydropsyc Hydropsyc Coleoptera Elimidae Other	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Margin: Predominant Org Other Common ( Density:	ganism: Organisr High High ganism: Organisr High High High ganism: Organisr High	ns:N Ns:N ns:N ns:N	Aoder Aoder Aoder Aoder Aoder	rate ate rate rate rate rate	Low Low Low Low Low	, , , ,	V≃ Very Overall An / / / / / / / / / / / / /	Abundant; A= Abundan iount (V=>1) Porifera, Bryoza Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera Hydropsyc Hydropsyc Coleoptera Elimidae Other	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Margin: Predominant Org Other Common ( Density: Diversity:	ganism: Organisr High High ganism: Organisr High High High ganism: Organisr High High	ns:N N N N N N	Aoder Aoder Aoder Aoder Aoder	rate ate rate rate rate rate rate	Low Low Low Low Low Low	, , , ,	V≃ Very Overall An / / / / / / / / / / /	Abundant; A= Abundan iount (V=>12 Porifera, Bryozz Turbellaria, Oli Isopoda, Amph Decapoda, Hyd Ephemeroptera Baetidae Heptageniic Other Zygoptera, Anis Plecoptera Hemiptera Megaloptera, N Trichoptera Hydropsyc Hydropsyli Other Coleoptera Elimidae Other Diptera	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina dae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Margin: Predominant Org Other Common ( Density: Diversity: Margin: Predominant Org Other Common ( Density: Diversity: Diversity:	ganism: Organisr High High ganism: Organisr High High High ganism: Organisr High High High	ns:N N N N N 	Aoder Aoder Aoder Aoder Aoder Aoder	rate rate rate rate rate rate rate	Low Low Low Low Low Low Low	, , , ,	V≃ Very Overall An / / / / / / / / / /	Abundant; A= Abundan 1000000000000000000000000000000000000	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae
Riffle: Predominant Org Other Common ( Density: Diversity: Run: Predominant Org Other Common ( Density: Diversity: Pool: Predominant Org Other Common ( Density: Diversity: Margin: Predominant Org Other Common ( Density: Diversity: Margin: Predominant Org Other Common ( Density: Diversity: Diversity: Other Common ( Density: Diversity: Diversity: Other Common ( Density: Diversity:	ganism: Organisr High High ganism: Organisr High High ganism: Organisr High ganism: Organisr High High	ns:N N N N N N	Aoder Aoder Aoder Aoder Aoder Aoder	rate rate rate rate rate rate rate	Low Low Low Low Low Low Low	, , ,	V= Very Overall An / / / / / / / / / / / / /	Abundant; A= Abundan 1000000000000000000000000000000000000	nt; C= Common; R= Rare 51; A= 150-101; C= 100-11; R= 10-1) oa gochaeta, Hirudinea ipoda racarina lae, Leptohyphidae, Caenidae soptera europtera chidae dae, Leptoceridae idae

**ChieEPA** 

Qualitative Habitat Evaluation Index and Use Assessment Field Sheet QHEI Score:

Stream & Location:	RM:	Date:	1	/
Scorers Full Name & Affiliation:	Northeast	- Ohio Regional S	ewer Dist	rict
River Code: STORET #: Lat./Long.:	/8	'	Office v	verified ocation
1] SUBSTRATE Check ONLY Two substrate TYPE BOXES; estimate % or note event type present	NE(0r2)	average)		
BEST TYPES POOL BIEFLE OTHER TYPES POOL BIEFLE ORIGIN		QUALI	TY	
		HEAVY [-	2]	
	SILT		[E[-1] ( [0]	Substrate
GRAVEL [7]      GRAVEL [7]      HARDPAN [0]		FREE [1]		
	SEDDEON		VE [-2] TE [-1]	
NUMBER OF BEST TYPES: 4 or more [2] sludge from point-sources)	Ē		[0]	Maximum 20
Comments		NONE [1]		
2] INSTREAM COVER Indicate presence 0 to 3: 0-Absent; 1-Very small amounts or if more commo	n of margin	al AMOU	JNT	
quality; 3-Highest quality in moderate or greater amounts (e.g., very large boulders in deep of fast water	, large	Check ONE (O	2 & aver	age)
UNDERCUT BANKS [1] POOLS > 70cm [2] OXBOWS. BACKWATE	pools. RS [1]		>75% [11] 25-75% [7	1 71
OVERHANGING VEGETATION [1] ROOTWADS [1] AQUATIC MACROPHY	TES [1]	SPARSE 5-4	25% [3]	
SHALLOWS (IN SLOW WATER) [1]BOULDERS [1]LOGS OR WOODY DEE ROOTMATS [1]	BRIS [1]		SENT <5%	6[1]
Comments		٨	Cover Maximum	
			20	$\Box$
3] CHANNEL MORPHOLOGY Check ONE in each category (Or 2 & average)				
SINUOSITY DEVELOPMENT CHANNELIZATION STABILITY				
			Channel	
Comments		Λ	Maximum	
			20	
4] BANK EROSION AND RIPARIAN ZONE Check ONE in each category for EACH BANK (O.	r 2 per bani	k & average)		
	ΗА			E 741
NONE / LITTLE [3] I MODERATE 10-50m [3] I SHRUB OR OLD FIELD [2]		URBAN OR IND	USTRIAL	. [0]
	[1]	MINING / CONS	TRUCTIO	N [0]
	Indicat past 1	e predominant la 00m riparian.	nd use(s) Rinarian	$\square$
Comments		٨	Aaximum	
			10	
5] POOL / GLIDE AND RIFFLE / RUN QUALITY MAXIMUM DEPTH CHANNEL WIDTH CURRENT VELOCITY		Recreation	Potenti	ial
Check ONE (ONLY!) Check ONE (Or 2 & average) Check ALL that apply		Primary	Contac	t
> 1m [6] POOL WIDTH > RIFFLE WIDTH [2] TORRENTIAL [-1] SLOW [1]		Secondary	y Conta	ct
0.4<0.7m [2] POOL WIDTH < RIFFLE WIDTH [0] FAST [1] INTERMIT	TENT [-2]	Circle one and co	mment on ba	ck)
0.2<0.4m [1]     MODERATE [1]     DEDDIES [1     Indicate for reach - pools and ri	] fflos		Pool / Current	$\bigcirc$
Comments	mes.	/	Maximum	
Indicate for functional rifflag. Dast grass must be large anough to support			12	
of riffle-obligate species: Check ONE (Or 2 & average).	a popula		RIFFLE (m	netric=0]
RIFFLE DEPTH RUN DEPTH RIFFLE / RUN SUBSTRATE RIFI	FLE / RU	N EMBEDDE	DNESS	5
BEST AREAS > 10cm [2] MAXIMUM > 50cm [2] STABLE (e.g., Cobble, Boulder) [2]		IONE [2]		
BEST AREAS < 5cm UNSTABLE (e.g., Fine Gravel, Sand) [0]		ODERATE [0]	Riffle /	$\square$
Comments		XTENSIVE [-1]	Maximum	$\cup$
	0/ 01 15		8	T
DRAINAGE AREA DI MODERATE [6-10]	%GLID		Gradient Maximum	
( mi <sup>2</sup> ) 🗆 HIGH - VERY HIGH [10-6] %RUN:	%RIFFL	E:( /	10	
EPA 4520			06/	16/06

UREMENTS width depth ax. depth ax. <sup>2</sup> width ratio e:	
F] MEAS x width x width max. depth max. depth x bankfull bankfull x bankfull r bankfull r hodpron entrench. Legacy Tre	
<b>E</b> <i>J</i> <b>ISSUES</b> WWTP / CSO / NPDES / INDUSTRY HARDENED / URBAN / DIRT&GRIME CONTAMINATED / LANDFILL BMPs-CONSTRUCTION-SEDIMENT LOGGING / IRRIGATION / COOLING BANK / EROSION / SURFACE FALSE BANK / MANURE / LAGOON WASH H <sub>2</sub> 0 / TILE / H <sub>2</sub> 0 TABLE ACID / MINE / QUARRY / FLOW NATURAL / WETLAND / STAGNANT PARK / GOLF / LAWN / HOME ATMOSPHERE / DATA PAUCITY	
Circle some & COMMENT	
<b>DJ MAINTENANCE</b> PUBLIC / PRIVATE / BOTH / NA ACTIVE / HISTORIC / BOTH / NA YOUNG-SUCCESSION-OLD SPRAY / SNAG / REMOVED MODIFIED / DIPPED OUT / NA LEVEED / DIPPED OUT / NA LEVEED / ONE SIDED RELOCATED / CUTOFFS MOVING-BEDLOAD-STABLE ARMOURED / SCOURED IMPOUNDED / DESICCATED FLOOD CONTROL / DRAINAGE	
BJAESTHETICS         BJAESTHETICS         NUISANCE ALGAE         INVASIVE MACROPHYTES         EXCESS TURBIDITY         DISCOLORATION         EXCESS TURBIDITY         INVASIVE DORATION         INUSANCE ODOR         INUSANCE ODOR         SLUDGE DEPOSITS         CSOS/SSOS/OUTFALLS         ATION       AREA DEPTH	
0.5 Km 0.2 Km 0.15 Km 0.15 Km 0.15 Km 0.15 Km 0.12	tream Drawing:
	0.3 Km       CLARITY       B) AESTHETICS       D) MAINTENANCE       Cite some & COMMENT       E) ISSUES       F) MEASUES       F) ME

**ChieEPA** Primary Headwater Habitat Evaluation Form

HHEI Score (sum of metrics 1, 2, 3) :

1

SITE NUMBERRIVER B	SIN DRAINAGE AREA (mi²)
LENGTH OF STREAM REACH (ft) LAT LO	IG RIVER CODE RIVER MILE
NOTE: Complete All Items On This Form - Refer to "Field Ev	uation Manual for Ohio's PHWH Streams" for Instructions
STREAM CHANNEL NONE / NATURAL CHANNEL	COVERED RECOVERING RECENT OR NO RECOVERY
1.       SUBSTRATE (Estimate percent of every type of substrate predimension of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of significant substrate types found (Max of 32). Add total number of signit (Max of 32). Add total number of significan	ent. Check ONLY two predominant substrate TYPE boxes Max of 8). Final metric score is sum of boxes A & B. SILT [3 pt] LEAF PACK/WOODY DEBRIS [3 pts] FINE DETRITUS [3 pts] CLAY or HARDPAN [0 pt] MUCK [0 pts] ARTIFICIAL [3 pts] (B) (B)
Bldr Slabs, Boulder, Cobble, Bedrock (A) SCORE OF TWO MOST PREDOMINATE SUBSTRATE TYPES:	Sates we Parcentage 0%     (B)     A + B       TOTAL NUMBER OF SUBSTRATE TYPES:     1
2. Maximum Pool Depth (Measure the maximum pool depth with evaluation. Avoid plunge pools from road culverts or storm water	in the 61 meter (200 ft) evaluation reach at the time of ipes) (Check ONLY one box): Max = 30
<ul> <li>&gt; 30 centimeters [20 pts]</li> <li>&gt; 22.5 - 30 cm [30 pts]</li> <li>&gt; 10 - 22.5 cm [25 pts]</li> </ul>	> 5 cm - 10 cm [15 pts] < 5 cm [5 pts] NO WATER OR MOIST CHANNEL [0 pts]
COMMENTS	MAXIMUM POOL DEPTH (centimeters):
BANK FULL WIDTH (Measured as the average of 3-4 measur           > 4.0 meters (> 13') [30 pts]           > 3.0 m - 4.0 m (> 9' 7" - 13') [25 pts]           > 1.5 m - 3.0 m (> 9' 7" - 4' 8") [20 pts]	ments)         (Check ONLY one box):         Bankfull           > 1.0 m - 1.5 m (> 3' 3" - 4' 8") [15 pts]         Width           ≤ 1.0 m (<=3' 3") [5 pts]
COMMENTS	AVERAGE BANKFULL WIDTH (meters):
This information         RIPARIAN ZONE AND FLOODPLAIN QUALITY         RIPARIAN WIDTH       FLOODPLAIN QUAL         L       R       (Per Bank)       L       R       (Most Pred         Wide >10m       Immature For       Immature For       Immature Field         Moderate 5-10m       Immature Field       Residential         None       Fenced Pa         COMMENTS       Flow REGIME (At Time of Evaluation)       (Check ONLY of Stream Flowing         Subsurface flow with isolated pools (Interstitial)       COMMENTS         SINUOSITY (Number of bends per 61 m (200 ft) of channe       1.0         0.5       1.5         STREAM GRADIENT ESTIMATE	1 must also be completed         DTE: River Left (L) and Right (R) as looking downstream ☆         Y         minant per Bank)       L R         st, Wetland       Conservation Tillage         rest, Shrub or Old       Urban or Industrial         Park, New Field       Open Pasture, Row Crop         ure       Mining or Construction         He box):       Moist Channel, isolated pools, no flow (Intermittent)         Dry channel, no water (Ephemeral)         H)       (Check ONLY one box):         2.0       3.0         2.5       3.0
Flat (0,5 ft/100 ft) Flat to Moderate Moderate (2 ft/100 f	Moderate to Severe Severe (10 ft/100 ft)

October 24, 2002 Revision

ADDITIONAL STREAM INFORMATION (This Information Must Also be Completed):	
QHEI PERFORMED? - Yes No QHEI Score (If Yes, At	tach Completed QHEI Form)
DOWNSTREAM DESIGNATED USE(S)	
WWH Name:	Distance from Evaluated Stream
CWH Name:	Distance from Evaluated Stream
EWH Name:	Distance from Evaluated Stream
MAPPING: ATTACH COPIES OF MAPS, INCLUDING THE ENTIRE WATERSHE	DAREA. CLEARLY MARK THE SITE LOCATION
USGS Quadrangle Name: NRCS Soil Map	Page: NRCS Soil Map Stream Order
County: Wyandot Township / City:	
MISCELLANEOUS	
Base Flow Conditions? (Y/N):_Y Date of last precipitation:	Quantity: 0.00
Photograph Information:	
Elevated Turbidity? (Y/N): Y Canopy (% open): 0%	
Were samples collected for water chemistry? (Y/N):	and attach results) Lab Number:
Field Measures: Temp (*C) Dissolved Oxygen (mg/l) pH (S.U.)	Conductivity (µmhos/cm)
Is the sampling reach representative of the stream (Y/N) If not, please explain:	
Additional comments/description of pollution impacts:	
Performed? (Y/N): ()f Yes, Record all observations. Voucher collections option	al. NOTE: all voucher samples must be labeled with the site
ID number. Include appropriate field data sheets from the P	rimary Headwater Habitat Assessment Manual)
Fish Observed? (Y/N) Voucher? (Y/N) Salamanders Observed? (Y/N)	Voucher? (Y/N)
Frogs or Ladpoles Observed? (Y/N) Y Voucher? (Y/N) Y Aquatic Macroinvertebr	ates Observed? (Y/N) Y Voucher? (Y/N)
Comments Regarding Biology:	

### DRAWING AND NARRATIVE DESCRIPTION OF STREAM REACH (This must be completed):

Include important landmarks and other features of interest for site evaluation and a narrative description of the stream's location





Lake / Lacustuary (Lentic) QHEI Field Sheet Ohio	onmentai clion Agency QHEI Score:
RIVERCODE RIVERMILE WATERBODY	DISTANCE ASSESSED (m):
1] SUBSTRATE (Check ONLY Two Substrate TYPE BOXES; Estimate % or note every type p         TYPE       SHORE BOTTOM       SHORE BOTTOM         Check ONLY Two Substrate TYPE BOXES; Estimate % or note every type p         TYPE       SHORE BOTTOM       SUBSTRATE ORIGN         Check ONLY Two Substrate TYPE BOXES; Estimate % or note every type p       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [10]       Check ONE (or 2 & AVERAGE         Check ONE [11]       Check ONE (or 2 & AVERAGE         Check ONE [12]       Check ONE (or 2 & AVERAGE         Check ONE [13]       Check ONE (or 2 & AVERAGE         Check ONE [14]       Check ONE [15]         Check ONE [15]       Check ONE [16]         Check ONE [16]       Check ONE [16]         Check ONE [16]       Check ONE [16]	SLT: D-SLT INOUTRAL [1] D-CLAY [2] SLT: D-SLT INOUTRAL [1] D-SLT INOUTRAL [1] D-SL
COMMENTS:	
2] COVER TYPES     TYPE: (Check All That Apply]     AM       D-OFF-SHORE SAND BARS [4]     D-DEEPWATER>1 M[1]     D-WETLAND POOLS [1]     D-H       D-OVERHANGING VEGETATION [1]     D-ROOTWADS [1]     D-SUBMERGED AQUATIC VEG. [4]     D-H       D-SHALLOWS (ON BEACH) [1]     D-BOULDERS [1]     D-LOGS OR WOODY DEBRIS [1]     D-H       D-ROOTMATS [1]     D-SAND BEACH [1]     D-GRAVEL BEACH [1]     D-H       COMMENTS:     D-SAND BEACH [1]     D-GRAVEL BEACH [1]     D-H	QUNT:         (Check ONLY One or check2 and AVERAGE)           EXTENSIVE > 75% [9]         Cover           MODERATE 25-75% [7]         Cover           SPARSE 5-25% [3]         Max 20
SHORE SINUOSITY       DEVELOPMENT         D-HIGH [2]       D-EXCELLENT [5]         D-MODERATE [4]       D-EXCELLENT [5]         D-LOW [3]       D-FAIR [3]         D-HONE [1]       D-FOOR [1]         SHORE to BOTTOM SLOPE MORPHOLOGIES       AVERAGE DEPTH (of 5 measures)         D-SLOPE < 15 deg. [0]	-CEMENTED [-1] -CEMENTED [-1] -REP RAPPED [1] -REI RAADS [2] -REI RAAD TIES [-1] -DREDGED [-1] -TWO SIDE CHANNEL -TWO SIDE CHANNEL
COMMENTS	Max 20
4] RIPARIAN ZONE AND BANK EROSION (Check OVE box PER bank or 2 and AVERAGE)	Shore Right Looking East or South on Lake 🔶 🔶
RIPARIAN WIDTH         SHORE LINE QUALITY (PAST 100 FOOT RIPARIAN)           L R (PerBank)         L R (Most Redominant PerBank)	Shore Right Looking Toward Lake in Lacustuary 🛧 BANK EROSION L R (PerBank) Bristian
Di Di-Mude > 50 m [4]       Di Di-FOREST, WETLAND, LAKE [3]       Di Di-CONSERVATION TILL         Di Di-Moderate 10-50 m [3]       Di Di-SHRUB OR OLD FIELD [2]       Di Di-URBAN OR INDUSTR         Di Di-NARROW 5-10 m [2]       Di Di-VINEYARD, ORCHARD [2]       Di Di-OPENPASUTRE, RO         Di Di-VERY NARROW < 5 m [1]	LIAGE [1] BAL [0] WCROP [0] TION [0]
COMMENTS	
5) AQUATIC VEGETATION QUALITY: <u>PLANT SPECIES OBSERVED</u> (Sum All Scores) (Score all for observed abundance: ABUNDANT = [3]; COMMON = [5]; FEW= [1]; UNCOMMON = [0])	
-Pond Lilles (NYMPHAEA)	A)Wild Rice (ZIZANIA) Vegezion
(Score all for observed abundance: ABUNDANT = [-2]; COMMON = [-1]; FEW = [0])	
-Purple Loosestrife -Reed Grass -Eurasian Milfoli -Cattalis -Alga	e (mats)Algae (planktonic)

June 2010

is the Sampling Reach Representa	live of Area	Habitat? (Y/N) if Not,	Explain:			
Depth measures:     Zebra Mussel/Quagga Mussel Cov	erage	 _↓>60%60->25%	□-25->10% □-<10	<u>-&gt;1% □</u> -1-0%		
First Sampling Pass: Second Sampling Pass: Third Sampling Pass:	Gear	Distance	Water Clarity	Wave Heighi	Subjective Reling (1-10) Photos:	Aesthetic Rating (1-10)
WATERBODY MEASUREMENTS	AVE		AVERAGE DEPTH:_	Mexin	num Depth:	)

June 2010

|--|

### PHWH STREAM BIOLOGICAL CHARATERISTICS FIELD SHEET:

1. Fish: Voucher Speci Sample Method No Evaluation	imens Retained?-(select)	t) Yes Time Spent (minutes): Stream Length Assessed (meters)
Species	Number Caught	Notes
Blank	0	
	0	
	0	
	0	
	0	

### 

Species (Genus)	# Larvae	# Juveniles/Adults	Total Number
<b>Mountain Dusky</b> (Desmognathus ochrophaeus)	0	0	0
Northern Dusky (Desmognathus fuscus)	0	0	0
<b>Two-lined</b> (Eurycea bislineata)	0	0	0
Long-tailed (Eurycea longicauda)	0	0	0
<b>Cave</b> (Eurycea lucifuga)	0	0	0
<b>Red</b> (Pseudotriton ruber)	0	0	0
Mud (Pseudotriton montanus)	0	0	0
<b>Spring</b> (Gyrinophilus porphyriticus)	0	0	0
Mole spp. (Ambystoma spp.)	0	0	0
Four-toed (Hemidactylium scutatum)	0	0	0
Other (name)	0	0	0
Total	0	0	0

Notes on Vertebrates:

4/1/03

PHWH FORM - Page 3

### 3. Macroinvertebrate Scoring Sheet:

THE HEADWATER MACROINVERTEBRATE FIELD EVALUATION INDEX (HMFEI) SCORING SHEET

Indicate Abundance of Each Taxa Above each White Box.

Record HMFEI Scoring Value Points Within each Box.

For EPT taxa, also indicate the different taxa present.

Key: $V = V$	Very Abundant	(>50); A = Abunda	ant (10	-50); C	= Common ( 3 -9);	$\mathbf{R} = \text{Rare}$	(<3)
Sessile Animals (Pori	fera,	Crayfish ( <b>Decapoda</b> )			Fishfly Larvae		
Cnidaria, Bryozoa)			NA		(Corydalidae)	NA	
(HMFEI pts = 1)		(HMFEI pts = 2)	NA	U	(HMFEI pts = 3)	NA	
Aquatic Worms (Turl	bellaria, Hirudinea,	Dragonfly Nymphs			Water Penny Beetles		
Oligochaeta)	NA O	(Anisoptera)	ΝΔ	0	(Psephenidae)	NA	
(HMFEI pts = 1)		(HMFEI pts = 2)			(HMFEI pts = 3)		
Sow Bugs		Riffle Beetles (Dryopidae,			Cranefly Larvae		
(Isopoda)		Elmidae, Ptilodactylidae)	NA	0	(Tipulidae)	NA	
(HMFEI pts = 1)		(HMFEI pts = 2)			(HMFEI pts = 3)		
Scuds (Amphipoda)		Larvae of other Flies (enter	name in c	omments)	EPT	ТАХА*	
(HMFEI pts = 1)		(Diptera):				0	
		(HMFEI pts = 1)	NA	U	Total No. EPT Taxa =		
Water Mites (Hydrac	arina)	Midges (Chironomidae)			Mayfly Nymphs (Ephen	neroptera)	
(HMFEI pts = 1)		(HMFEI pts = 1)			Taxa Present:	0	
	NA O		AL A		[HMFEI pts =		
			NA	<u> </u>	No. Taxa (x) 3]	NA	0
Damselfly Nymphs		Snails					
(Zygoptera)		(Gastropoda)	NA				
(HMFEI pts = 1)		(HMFEI pts = 1)					
Alderfly Larvae		Clams			Stonefly Nymphs (Pleco	optera)	
(Sialidae)		(Bivalvia)			Taxa Present:	0	
(HMFEI pts = 1)		(HMFEI pts = 1)			HMFEI pts =	NA	
			NA	0	No. Taxa (x) 3]	NA	U
Other Beetles		Other Taxa :					
(Coleoptera)							
(HMFEI pts = 1)							
Other Taxa:		Other Taxa:			Caddisfly Larvae (Trich	ioptera)	
					Taxa Present:	0	
L					HMFEI pts =	NIA	
					No. Taxa (x) 3]	NA	
Other Taxa:		Other Taxa					
				DT 1 CC			1 6
			*Note: E	P1 identifi	cation based upon Family	or Genus leve	l of taxonom
Voucher Sample ID_			Time S	pent (minu	tes):		
Notes on Macroinver	tebrates: (Predomir	ant Organisms; Other Com	non Organ	isms; Diver	rsity Estimate)		
						<b></b>	
	Final HMFE	EI Calculated Scor	e (Sum	of All	White Box Score	es) =	

IF Final HMFEI Score is > 19, Then CLASS III PHWH STREAM IF Final HMFEI Score is 7 to 19, Then CLASS II PHWH STREAM

IF Final HMFEI Score is < 7, Then CLASS I PHWH STREAM

Inar HWI EI SCOLE IS < 7, THEIL CEASS TO THIWH STREAM

PHWH FORM - Page 4

Reset Form

4/1/03

Stream:			Date:		Co	ollectors:		
Gage Statio	on and ID:	• • • • • •			_Daily Mean	Discharge:		ft³/sec
Was this sar	mple taken	during or foll	lowing a wet	weather e	vent?	YES / 1	NO	
Water Quali	ity Meters	Used:						
Time (hrs):			River I	Mile (Site)	:			
<u>Weather:</u> Ste	Clear eady Rain	Partly Cloud Heavy S	y Overca Snow Melt	ast Lig Oth	ght Rain/Show	vers He	eavy Rain	
<u>Flow:</u> Dr	ry Inte	rmittent	Minimal	Baselin	e/Normal	Elevated	Flood	
HD Status:		OK	Other:					
<u>Color:</u>	Clear	Mu	ddy	Tea	Milky	Ot	her:	
<u>Odor:</u>	Normal	Petroleum	Anae	robic	Sewage	Chemical	Other:	
Surface Coa	<u>ating:</u>	None	Foam	Oily	Scum	Other:		
Field Param	neters:	Conductivity	y (µmhos/cm	i):		Sp. Cond. (µ	mhos/cm):	
		Dissolved O	xygen (mg/L	.):		D.0	). (%):	
		Ten	nperature (°C	:):		pH	(s.u.):	
Turbidity	y 1 (NTU):	Ten	nperature (°C Turbidit	:): y 2 (NTU)	:	pH Avera	(s.u.): lge (NTU):	
Turbidity General Cor	y 1 (NTU): mments:	Ten	nperature (°C 	c): y 2 (NTU)	:	pH Avera	(s.u.): ge (NTU):	
Turbidity General Con	y 1 (NTU): mments:	Ten	nperature (°C Turbidit	c): y 2 (NTU)		pH Avera	(s.u.): ge (NTU):	
Turbidity General Con	y 1 (NTU): mments:	Ten	nperature (°C Turbidit	;): y 2 (NTU)		pH Avera	(s.u.): ge (NTU):	
Turbidity General Cor  Fime (hrs):	y 1 (NTU): mments:	Ten	nperature (°C Turbidit	S): Y 2 (NTU) Mile (Site)	):	pH Avera	(s.u.): lge (NTU):	
Turbidity General Con  Fime (hrs): <u>Weather:</u> Sto	y 1 (NTU): mments: Clear eady Rain	Partly Cloud Heavy S	nperature (°C Turbidit  River 1 ly Overca Snow Melt	:): y 2 (NTU) Mile (Site) ast Lig Otl	): ): ght Rain/Show her:	pH Avera	(s.u.): ge (NTU): eavy Rain	
Turbidity General Con  Fime (hrs): Weather: Sto Flow: Dr	y 1 (NTU): mments: Clear eady Rain ry Inte	Ten Partly Cloud Heavy S	nperature (°C Turbidit River I y Overca Snow Melt Minimal	:): y 2 (NTU) Mile (Site) ast Lig Oth Baselin	): ): ght Rain/Shov her: e/Normal	pH Avera  vers Ho Elevated	(s.u.): ge (NTU): eavy Rain Flood	
Turbidity General Con Time (hrs): Weather: Sta Flow: Dr HD Status:	y 1 (NTU): mments: Clear eady Rain ry Inte	Partly Cloud Heavy S ermittent OK	nperature (°C Turbidity  River I dy Overca Snow Melt Minimal Other:	c): y 2 (NTU) Mile (Site) ast Lig Otl Baselin	): ): ght Rain/Shov her: e/Normal	pH Avera	(s.u.): ge (NTU): eavy Rain Flood	
Turbidity General Con Fime (hrs): Weather: Sto Flow: Dr HD Status: Color:	y 1 (NTU): omments: Clear ready Rain ry Inte Clear	Partly Cloud Heavy S ermittent OK Mu	nperature (°C Turbidity  River I y Overca Snow Melt Minimal Other: uddy	:): y 2 (NTU) Mile (Site) ast Lig Oth Baselin Tea	: ght Rain/Shov her: e/Normal Milky	pH Avera  vers Ho Elevated	(s.u.): ge (NTU): eavy Rain Flood ther:	
Turbidity General Con Fime (hrs): Weather: Sto Flow: Dr HD Status: Color: Odor:	y 1 (NTU): omments: Clear ceady Rain ry Inte Clear Normal	Partly Cloud Heavy S crmittent OK Mu Petroleum	nperature (°C Turbidity  River I y Overca Snow Melt Minimal Other: Iddy Anae	c): y 2 (NTU) Mile (Site) ast Lig Baselin Tea crobic	: ght Rain/Show her: e/Normal Milky Sewage	pH Avera vers Ho Elevated Of Chemical	(s.u.): ge (NTU): eavy Rain Flood ther: Other:	
Turbidity General Con Fime (hrs): Weather: Sto Flow: Dr HD Status: Color: Odor: Surface Coa	y 1 (NTU): omments: Clear ceady Rain ry Inte Clear Normal ating:	Ten Partly Cloud Heavy S ermittent OK Mu Petroleum None	nperature (°C Turbidity  River I dy Overca Snow Melt Minimal Other: Iddy Anae Foam	c): y 2 (NTU) Mile (Site) ast Lig Oth Baselin Tea crobic Oily	: ght Rain/Show her: e/Normal Milky Sewage Scum	pH Avera  vers He Elevated Of Chemical Other:	(s.u.): ge (NTU): eavy Rain Flood ther: Other:	
Turbidity General Con Time (hrs): Weather: Status: Flow: Dr HD Status: Color: Odor: Surface Coa Field Param	y 1 (NTU): omments: Clear eady Rain ry Inte Clear Normal <u>ating:</u> neters:	Ten Partly Cloud Heavy S ermittent OK Mu Petroleum None Conductivity	nperature (°C Turbidity River l y Overca Snow Melt Minimal Other: ddy Anae Foam y (µmhos/cm	c): y 2 (NTU) Mile (Site) ast Lig Baselin Tea crobic Oily	: ght Rain/Show her: e/Normal Milky Sewage Scum	pH Avera vers Ho Elevated Of Chemical Other: Sp. Cond. (µ	(s.u.): ge (NTU): eavy Rain Flood ther: Other: 	
Turbidity General Con Time (hrs): Weather: Sta Flow: Dr HD Status: Color: Odor: Surface Coa Field Param	y 1 (NTU): omments: Clear ry Inte Clear Normal ating: neters:	Partly Cloud Heavy S ermittent OK Mu Petroleum None Conductivity Dissolved O	nperature (°C Turbidity  River I y Overca Snow Melt Minimal Other: Iddy Foam y (µmhos/cm xygen (mg/I	c): y 2 (NTU) Mile (Site) ast Lig Baselin Tea crobic Oily n):	: ght Rain/Show her: e/Normal Milky Sewage Scum	pH Avera  vers Ho Elevated  Chemical  Of Chemical  Other: Sp. Cond. (µ D.C	(s.u.): ge (NTU): eavy Rain Flood ther: Other: 	
Turbidity General Con Fime (hrs): Weather: Sta Flow: Dr HD Status: Color: Odor: Surface Coa Field Param	y 1 (NTU): omments: Clear ry Inte Clear Normal <u>ating:</u> neters:	Ten Partly Cloud Heavy S ermittent OK Mu Petroleum None Conductivity Dissolved O Ter	nperature (°C Turbidity  River I dy Overca Snow Melt Minimal Other: ddy Anae Foam y (µmhos/cm xygen (mg/I nperature (°C	c): y 2 (NTU) Mile (Site) ast Lig Baselin Tea crobic Oily n): _): C):	: ght Rain/Show her: e/Normal Milky Sewage Scum	pH Avera vers Ho Elevated Of Chemical Other: Sp. Cond. (µ D.C pH	(s.u.): ge (NTU): eavy Rain Flood ther: Other: 	
Turbidity General Con Fime (hrs): Weather: Status: Flow: Dr HD Status: Color: Odor: Surface Coa Field Param	y 1 (NTU): omments: Clear eady Rain ry Inte Clear Normal ating: neters: y 1 (NTU):	Ten Partly Cloud Heavy S ermittent OK Mu Petroleum None Conductivity Dissolved O Ten	nperature (°C Turbidity  River I dy Overca Snow Melt Minimal Other: Iddy Foam y (µmhos/cm xygen (mg/L nperature (°C Turbidit	<pre>c):</pre>	: ght Rain/Show her: e/Normal Milky Sewage Scum  ):	pH Avera wers Ho Elevated Of Chemical Other: Sp. Cond. (µ D.C pH Avera	(s.u.): ge (NTU): eavy Rain Flood ther: Other: 	

Sample ID:

Sample ID:

Modified January 28, 2015

Appendix B

Doromotor	Additional Nome	Test	2014	2014
Parameter	Additional Name	Test	Minimum Detection Limit	Practical Quantitation Limit
Alkalinity	Alkalinity	EPA 310.2	1.6 mg/L	5.0 mg/L
Mercury	Hg	EPA 245.1	0.006 µg/L	0.050 µg/L
Ammonia <sup>1</sup>	NH3	EPA 350.1	0.002 mg/L	0.020 mg/L
Nitrite	NO2	SM 4500 NO <sub>2</sub> <sup>12</sup> B <sup>2</sup>	0.001 mg/L	0.020 mg/L
Nitrite + Nitrate	NO <sub>2</sub> + NO <sub>3</sub>	EPA 353.2	0.003 mg/L	0.020 mg/L
Total Kjeldahl Nitrogen	TKN	EPA 351.2	0.081 mg/L	0.500 mg/L
Dissolved Reactive Phosphorus	DRPhos	EPA 365.1	0.003 mg/L	0.010 mg/L
Total Phosphorus	Total-P	EPA 365.1	0.003 mg/L	0.010 mg/L
Chloride	Chloride by IC	EPA 300.0	1.00 mg/L	5.00 mg/L
Sulfate	Sulfate by IC	EPA 300.0	0.500 mg/L	5.00 mg/L
Silver	Ag	EPA 200.8	0.009 µg/L	0.500 μg/L
Aluminum	Al	EPA 200.8	0.504 µg/L	5.000 μg/L
Arsenic	As	EPA 200.8	0.320 μg/L	1.000 μg/L
Barium	Ва	EPA 200.8	0.033 µg/L	0.500 μg/L
Beryllium	Be	EPA 200.8	0.540 μg/L	0.500 µg/L
Calcium	Ca	EPA 200.8	16.90 μg/L	125.0 μg/L
Cadmium	Cd	EPA 200.8	0.034 µg/L	0.500 µg/L
Cobalt	Co	EPA 200.8	0.056 µg/L	0.500 μg/L
Chromium	Cr	EPA 200.8	0.049 µg/L	0.500 μg/L
Copper	Cu	EPA 200.8	0.073 μg/L	1.000 μg/L
Iron	Fe	EPA 200.8	0.495 µg/L	5.000 μg/L
Potassium	К	EPA 200.8	3.695 μg/L	125.0 μg/L
Magnesium	Mg	EPA 200.8	2.077 μg/L	125.0 μg/L
Manganese	Mn	EPA 200.8	0.057 μg/L	1.000 μg/L
Molybdenum	Mo	EPA 200.8	0.017 μg/L	0.500 μg/L
5odium	Na	EPA 200.8	13.910 µg/L	125.0 μg/L
Nickel	Ni	EPA 200.8	0.066 μg/L	2.000 μg/L
Lead	Pb	EPA 200.8	0.058 μg/L	0.500 μg/L
Antimony	5b	EPA 200.8	0.018 µg/L	0.500 µg/L
Selenium	Se	EPA 200.8	0.383 µg/L	2.500 μg/L
Tin	5n	EPA 200.8	0.018 μg/L	0.500 μg/L
Strontium	Sr	EPA 200.8	0.049 μg/L	0.500 µg/L
Titanium	Ti	EPA 200.8	0.071 µg/L	1.000 μg/L
Thallium	TI	EPA 200.8	0.007 µg/L	0.500 μg/L
Vanadium	v	EPA 200.8	0.238 µg/L	5.000 μg/L
Zinc	Zn	EPA 200.8	0.241 μg/L	5.000 µg/L
Total Metals	Total Metals (calc.)	EPA 200.8	μg/L =(Cr μg/L)+(Cu με	g/L)+(Ni μg/L)+(Zn μg/L)
Hardness	Hardness (calc.)	5M 2340 <sup>2</sup>	CaCO3 mg/L =(2.497*Ca	mg/L)+(4.118*Mg mg/L)
		EPA 1603	1 colony	
Escherichia coli	E. coli	Colilert QT		
		(SM 9223 B 20th	1 MPN	1 MPN
Chlorophyll a	Chloronhylla	EQ) EPA 445 0	0.03.ug/l	0.15 ug/i
Chemical Oxygen Demand	COD	EPA 410.4	0.49 mg/l	10 mg/l
Biological Oxygen Demand	BOD	ENA 5210 <sup>2</sup>	2 mg/l	10 mg/c
Total Solids	TS	5M 3540 P 2	1.0 mg/i	5.0 mg/l
Total Suspended Solids	TSS	SM 2540 D <sup>2</sup>	0.5 mg/l	1.0 mg/l
Total Dissolved Solids		5M 2540 C	1.0 mg/l	5.0 mg/L
Turbidity **	105	FPA 180 1	0 1 NTU	0.2 NTU
Field Parameter		Tect	(Value Re	ported in)
nH		EPA 150 1 2	lange up	.u.
Conductivity		5M 2510A 2	115	/cm
Specific Conductivity	-	5M 2510A	110	/cm
Dissolved Oxygen	DO	SM 4500 0 G 2	μ., m	g/l
Temperature	Temn	FDA 1701 1 2		ог С
Turbidity **	i crip	FPA 180 1	N	ти
	1	CI / 100.1		

<sup>1</sup> Listed MDL/PQL is for undistilled samples. Any samples that require distillation will have a MDL = 0.020 mg/L, PQL = 0.100 mg/L

<sup>2</sup> Standard Methods for the Examination of Water and Wastewater, Method approved by Standard Methods Committee, 1997. Editorial revisions, 2011.

\*\* Turbidity will either be completed in the field or at the laboratory.

Appendix C

### **YSI** Environmental





A rugged, cost-effective multiparameter handheld system designed for the field!

### YSI 556 Multiparameter System

### Versatile, multiparameter handheld instrument

Rugged and reliable, the YSI 556 MPS (Multiprobe System) combines the versatility of an easy-to-use, easy-to-read handheld unit with all the functionality of a multiparameter system.

- Simultaneously measures dissolved oxygen, pH, conductivity, temperature, and ORP
- Field-replaceable electrodes
- Compatible with EcoWatch' for Windows' data analysis software
- Stores over 49,000 data sets, time and date stamped, interval or manual logging
- Three-year warranty on the instrument; one-year on the probes
- GLP assisting, records calibration data in memory
- Available with 4, 10, and 20-m cable lengths
- IP-67, impact-resistant, waterproof case
- Easy-to-use, screw-on cap DO membranes
- RS-232 interface for PC connection

### **Options to Fit Your Applications!**

• Battery Options – The unit is powered by alkaline batteries or an optional rechargeable battery pack with quick-charge feature.

• Optional Barometer – Internal barometer can be user-calibrated and displayed along with other data, used in dissolved oxygen calibrations, and logged to memory for tracking changes in barometric pressure. (Choose 556-02)

• Optional Flow Cell - The 5083 flow cell can be used for ground water applications or anytime water is pumped for sampling.

• Carrying Case – The instrument comes standard with YSI 5061, a soft-sided carrying case with enough space for the 556, a 20-meter cable, and calibrating supplies. An optional 5080 hard-sided carrying case is also available.

• Confidence Solution<sup>•</sup> - Quality assurance ensured. Quickly check conductivity, pH, and ORP readings with one solution.

www.YSI.com/556



+1 937 767 7241 800 897 4151 (US) www.ysi.com

YSI Environmental +1 937 767 7241 Fax +1 937 767 9353 environmental@ysi.com

YSI Integrated Systems +1 508 748 0366 systems@ysi.com

SonTek/YS1 +1 858 546 8327 inquiry@sontek.com

YSI Gulf Coast +1 225 753 2650 environmental@ysi.com

AMJ Environmental +1 727 565 2201 amj@ysi.com

YSI Hydrodata (UK) +44 1462 673 581 europe@ysi.com

YSI Middle East (Bahrain) +973 1753 6222 halsalem@ysi.com

YSI (Hong Kong) Limited +852 2891 8154 hongkong@ysi.com

YSI (China) Limited +86 532 575 3636 beijing@ysi-china.com.

YSI Nanotech (Japan) +81 44 222 0009 nanotech@ysi.com

YSI India +91 989 122 0639 sham@ysi.com

YSI Australia +61 7 390 17223 acorbett@ysi.com

ISO	9001
ISO	14001

EcoWatch, Pure Data for a Healthy Planet, Confidence Solution and Who's Minding the Planet? are registered trademarks of YSI Incorporated. Windows is a registered trademark of Microsoft.

D2009 YSI Incorporated	
Printed in USA 0709 W11-06	
ČE	
~ ~	

Ŷ	5	Lincorporated
		Who's Ainding
		the Planet?"

5563 MPS Senso	r Specification	ns
Dissolved Oxygen (% saturation)	Sensor Type Range Accuracy whichever is g Resolution	Steady state polarographic 0 to 500% air saturation 0 to 200% air saturation, ± 2% of the reading or ±2% air saturation, reater; 200 to 500% air saturation, ± 6% of the reading 0.1% air saturation
Dissolved Oxygen (mg/L)	Sensor Type Range Accuracy Resolution	Steady state polarographic 0 to 50 mg/L 0 to 20 mg/L, ± 2% of the reading or ±0.2 mg/L, whichever is greater; 20 to 50 mg/L, ± 6% of the reading 0.01 mg/L
Temperature	Sensor Type Range Accuracy Resolution	YSI Temperature Precision" thermistor -5 to 45°C ± 0.15°C 0.1°C
Conductivity	Sensor Type Range Áccuracy ± 1.0% of readi Resolution	4-electrode cell with autoranging 0 to 200 mS/cm ± 0.5% of reading or ± 0.001 mS/cm; whichever is greater (4-meter cable ing or ± 0.001 mS/cm; whichever is greater (20-meter cable) 0.001 mS/cm to 0.1 mS/cm (range-dependent)
Salinity	Sensor Type Range Accuracy Resolution	Calculated from conductivity and temperature 0 to 70 ppt ± 1.0% of reading or ±0.1 ppt, whichever is greater 0.01 ppt
pH (optional)	Sensor Type Range Accuracy Resolution	Glass combination electrode 0 to 14 units ±0.2 units 0.01 units
ORP (optional)	Sensor Type Range Accuracy Resolution	Platinum button -999 to +999 mV ± 20 mV 0.1 mV
Total Dissolved Solids (TDS)	Sensor Type Range Resolution	Calculated from conductivity (variable constant, default 0.65) 0 to 100 g/L 4 digits
Barometer (optional)	Range Accuracy Resolution	500 to 800 mm Hg ± 3 mm Hg within ± 10°C temperature range from calibration point 0.1 mm Hg

### YSI 556 Instrument Specifications

Size	11.9 cm width x 22.9 cm leñth (4.7 in. x 9 in.)
Weight with batteries	2.1 lbs. (916 grams)
Power	4 alkaline C-cells; optional rechargeable pack
Cables	4-, 10-, and 20-m (13.1, 32.8, 65.6 ft.) lengths
Warranty	3-year instrument; 1-year probes and cables
Communication Port	RS-232 Serial
Data Logger	49,000 data sets, date and time stamp, manual or logging, with user-selectable intervals

### 556 Ordering Information (Order all items separately)

556-01	Instrument (with 5061 large, soft-sided carrying case)	
556-02	Instrument with barometer option (with 5061 carrying case)	Contraction of the second second
5563-4	4-m cable and DO/temp/conductivity	
5563-10	10-m cable and DO/temp/conductivity	
5563-20	20-m cable and DO/temp/conductivity	
5564	pH Probe for any 5563 cable	
5565	pH/ORP Probe for any 5563 cable	
6118	Rechargeable battery pack kit (includes battery, adapter, charger)	Carl Contraction of the second
614	Ultra clamp, C-clamp mount	The second secon
616	Charger, cigarette lighter	
4654	Tripod (small tripod for instrument)	
5060	Small carrying case, soft-sided (fits instrument and 4-m cable)	
5065	Form-fitted carrrier with shoulder strap	
5080	Small carrying case, hard-sided (fits instrument, 4-m cable, flow	Freedom in Sector
	cell, batteries, membrane kit, calibration bottles)	The second second second
5083	Flow cell	Service and the service of the service
5085	Hands-free harness	The 5080 carrying case with 556.
5580	Confidence Solution <sup>•</sup> (insure probe accuracy with a simple field-	5663 Lookla and 5003 flow call
	check for conductivity, pH, and ORP)	5565-4 capie, and 5665 now cell.





The YSI 600XL and 600XLM

### YSI 600XL and 600XLM Sondes

### Measure multiple parameters simultaneously

The YSI 600XL and YSI 600XLM compact sondes measure eleven parameters simultaneously:

Temperature Conductivity Specific Conductance Salinity Resistivity TDS pH ORP Depth or Level Rapid Pulse<sup>\*\*</sup> DO (% and mg/L)

### Connect with Data Collection Platforms

Either sonde can easily connect to the YSI 6200 DAS (Data Acquisition System), YSI EcoNet<sup>™</sup> or your own data collection platform, via SDI-12 for remote and real-time data acquisition applications.

### Economical Logging System

The YSI 600XLM is an economical logging system for long-term, *in situ* monitoring and profiling. It will log all parameters at programmable intervals and store 150,000 readings. At one-hour intervals, the instrument will log data for about 75 days utilizing its own power source. The 600XL can also be utilized in the same manner with user-supplied external power.

- Either sonde fits down 2-inch wells
- Horizontal measurements in very shallow waters
- Stirring-independent Rapid Pulse\* dissolved oxygen sensor
- Field-replaceable sensors
- Easily connects to data collection platforms
- Available with detachable cables to measure depth up to 200 feet
- Compatible with YSI 650 Multiparameter Display System
- Use with the YSI 5083 flow cell for groundwater applications



Economical, multiparameter sampling or logging in a compact sonde

### Sensor performance verified\*

The  $6820 \lor 2$  and  $6920 \lor 2$  sondes use sensor technology that was verified through the US EPA's Environmental Technology Verification Program (ETV). For information on which sensors were performance-verified, turn this sheet over and look for the ETV logo.



www.ysi.com



To order, or for more info, contact YSI Environmental.

### +1 937 767 7241 800 897 4151 (US) www.ysi.com

YSI Environmental +1 937 767 7241 Fax +1 937 767 9353 environmental@ysi.com

Endeco/YSI +1 508 748 0366 Fax +1 508 748 2543 systems@ysi.com

SonTek/YSI +1 858 546 8327 Fax +1 858 546 8150 inquiry@sontek.com

YSI Gulf Coast +1 225 753 2650 Fax +1 225 753 8669 environmental@ysi.com

YSI Hydrodata (UK) +44 1462 673 581 Fax +44 1462 673 582 europe@ysi.com

YSI Middle East (Bahrain) +973 1753 6222 Fax +973 1753 6333 halsalem@ysi.com

YSI (Hong Kong) Limited +852 2891 8154 Fax +852 2834 0034 hongkong@ysi.com

YSI (China) Limited +86 10 5203 9675 Fax +86 10 5203 9679 heijing@ysi-china.com

YSI Nanotech (Japan) +81 44 222 0009 Fax +81 44 221 1102 nanotech@ysi.com

ISO 9001 ISO 14001 Yolkun Sjunegu, Ohio Facefry

ROX and Rapid Pulse are trademarks and Fee Watch, Pure Data for a Healthy Planet and White Minding the Planet? are registered trademarks at VM Incorported.

© 2007 YSI Incorporated Printed in USA 0107 E55-01

Theorem shifts dwith the ETV lagranger advanted during TV pagareneous the 134 FUTCH to Laternature multiple type of parameter during the problem of the transmission of the transmission

Y S1 incorporated Who's Minding the Planet?

### YSI 600XL & 600XLM Sensor Specifications

	Ronge	Resolution	Accuracy	
Dissolved Oxygen % Saturation EIV 6562 Rapid Pulse" Sensor*	0 to 500%	0.1%	0 to 200%: ±2% of reading or 2% air saturation whichever is greater; 200 to 500%: ±6% of reading	
Dissolved Oxygen mg/L 6562 Rapid Pulse" Sensor*	0 to 50 mg/L	0.Q1 mg/L	0 to 20 mg/L: ± 0.2 mg/L or 2% of reading, whichever is greater; 20 to 50 mg/L: ±6% of reading	
Conductivity 6560 Sensor* ETV	0 to 100 mS/cm	0.001 to 0.1 mS/cm (range dependent)	±0.5% of reading + 0,001 m\$/cm	
Salinity	0 to 70 ppt	0.01 ppt	±1% of reading or 0.1 ppt, whichever is greater	
Tempérolure 6560 Sensor* ETV	-5 tó +50°C	0.01°C	±0.15°C	
pH 6561 Sensor* ETV	0 to 14 units	0.01 init	±0.2 unit	
ORP	-999 to +999 mV	0.1 mV	±20 mV	
D <b>epth &amp; Leve</b> l Medium Shallow Vented Level	0 to 200 ft, 61 m 0 to 30 ft, 9.1 m 0 to 30 ft, 9.1 m	0.001 ft, 0.001 m 0.001 ft, 0.001 m 0.001 ft, 0,001 m	±0,4 ft,±0.12 m ±0,06 ft,±0.02 m ±0,01 ft, 0.003 m	

 Report nutputs of specific conductance (conductivity corrected to 25° C), reastivity, and total dissolved solids are also provided. These values are automatically calculated (rum conductivity according to algorithms found in Standard Methods for the Examination of Water and Wastewater (ed 1989)

YSI 600XL & 600XLM Sonde Specifications				
Medium		Fresh, set or polluted water		
Temperature	Operating Storage	-5 to +50°C -10 to +60°C		
Communications		RS-232, SDI-12		
Software		EcoWatch*		
Dimensions 400401 400404	Diameter Langth Weight	1.65 in, 4.19 cm   1.65 in, 4.9 cm 16 in, 40.6 cm   21.3 in, 54.1 cm 1.3 lbs, 0.59 kg   1.5 lbs, 0.69 kg		
Power Internal	External 600XLM only)	12 V DC 4 AA-size alkaline batteries		





### HI 98129 Combo pH/EC/TDS/Temperature Tester with Low Range EC



### Description

The HI 98129 Combo waterproof tester offer high accuracy pH, EC/TDS and temperature measurements in a single tester! No more switching between meters for your routine measurements. The waterproof Combo (it even floats) has a large easy-to-read, dual-level LCD and automatic shut-off. pH and EC/TDS readings are automatically compensated for the effects of temperature (ATC). This technologically advanced tester has a replaceable pH electrode cartridge with an extendable cloth junction as well as an EC/TDS graphite electrode that resists contamination by salts and other substances. This gives these meters a greatly extended life. Your tester no longer needs to be thrown away when the pH sensor is exhausted.

The EC/TDS conversion factor is user selectable as is the temperature compensation coefficient (ß). Fast, efficient, accurate and portable, the Combo pH, EC/TDS and temperature tester brings you all the features you've asked for and morel

Specifications

Range	pН	0.00 to 14.00 pH
Range	EC	0 to 3999 µS/cm
Range	TDS	0 to 2000 ppm
Range	Temperature	0.0 to 60.0°C / 32 to 140.0°F
Resolution	pН	0.01 pH
Resolution	EC	1 µS/cm
Resolution	TDS	1 ppm
Resolution	Temperature	0.1°C / 0.1°F
Accuracy	pН	±0.05 pH
Accuracy	EC/TDS	±2% F.S.
Accuracy	Temperature	±0.5°C / ±1°F
Temperature		pH: automatic; EC/TDS: automatic with ß adjustable
Compensation		from 0.0 to 2.4% / °C
Calibration	pH	automatic, 1 or 2 points with 2 sets of memorized
		buffers
		(pH 4.01 / 7.01 / 10.01 or 4.01 / 6.86 / 9.18)
Calibration	EC/TDS	automatic, 1 point
<b>TDS Conversion Facto</b>	pr	adjustable from 0.45 to 1.00
pH Electrode		HI 73127 (replaceable; included)
Environment		0 to 50°C (32 to 122°F); RH max 100%
Battery Type / Life		4 x 1.5V / approx. 100 hours of continuous use;
		auto-off after 8 minutes of non-use
Dimensions		163 x 40 x 26 mm (6.4 x 1.6 x 1.0")
Weight		100 g (3.5 oz.)

HQ30d Portable pH, Conductivity, Dissolved Oxygen (DO), ORP, and... http://www.hach.com/hq30d-portable-ph-conductivity-dissolved-oxyg...



### 2100P and 2100P IS Portable Turbidimeter

### Features and Benefits

### Laboratory Quality in a Portable Unit

The Hach 2100P and 2100P IS Portable Turbidimeters offer a level of performance previously possible only with laboratory instruments. Microprocessor-controlled operation and Hach's unique Ratio<sup>™</sup> optics bring great accuracy, sensitivity, and reliability to field and in-plant testing.

### **Two Models for Specific Requirements**

- 2100P Turbidimeter—Get fast, accurate turbidity testing in the field or the lab, over a wide range of samples. Compliant with USEPA Method 180.1 design criteria.
- 2100P IS Turbidimeter—Designed to meet international standards that mandate measurement using an LED light source.

### **Two-detector Optical System**

The two-detector optical system compensates for color in the sample, light fluctuation, and stray light, enabling analysts to achieve laboratory-grade performance on a wide range of samples, even under difficult, onsite conditions.



The Hach 2100P and 2100P IS Portable Turbidimeters bring laboratory-level performance on-site, offering fast, accurate results and the ease-of-use analysts demand in the field. With a measurement range of 0 to 1000 NTU and a resolution of 0.01 NTU, the 2100P turbidimeter is ideal for regulatory monitoring, process control or field studies.

W
P

D

### Specifications\*

	2100P	2100P IS			
Measurement Method	Nephelometric Ratio				
Regulatory	Meets EPA Method 180.1 Meets EN ISO 7027				
Light Source	Tungsten lamp	Llght-emitting diode (LED) @ 860 nm			
Range					
Automatic Range Mode	0 to 1000 NTU	0 to 1000 FNU			
Manual Range Selection	0 to 9.99, 0 to 99.9 and 0 to 1000 NTU	0 to 9.99, 0 to 99.9 and 0 to 1000 FNU			
Accuracy	±2% of reading plus stray light				
Repeatability	±1% of reading, or 0.01 NTU, whichever is greater	±1% of reading, or 0.01 FNU, whichever is greater			
Resolution	0.01 on lowest range				
Signai Averaging	Selectable on/off				
Power Requirement	4 AA alkaline batteries or optional battery eliminator				
Battery Life, Typical	300 tests with signal average mode off				
	180 tests with signal average mode on				
Operating Temperature	0 to 50°C (32 to 122°F)				
Sample Required	15 mL (0.5 oz.)				
Sample Cells	60 x 25 mm (2.36 x 1 in.) borosilicate glass with screw	60 x 25 mm (2.36 x 1 in.) borosilicate glass with screw caps			
Dimensions	22.2 x 9.5 x 7.9 cm (8.75 x 3.75 x 3.12 in.)	22.2 x 9.5 x 7.9 cm (8.75 x 3.75 x 3.12 in.)			
Weight	0.5 kg (1.1 lb.); shipping weight 2.7 kg (6 lb.)				
Warranty	2 years				

\*Specifications subject to change without notice.

DW = drinking water WW = wastewater municipal PW = pure water / power IW = industrial water E = environmental C = collections FB = food and beverage



### 2100Q and 2100Q is Portable Turbidimeter



The Hach 2100Q and 2100Q is Portable Turbidimeters offer unsurpassed ease of use and accuracy in turbidity measurement. Only Hach offers this unique combination of advanced features, such as assisted calibration and simplified data transfer, and measurement innovation, giving you accurate results every time.



### **Easy Calibration and Verification**

Hach 2100Q and 2100Q *is* Portable Turbidimeters provide confidence your measurements are right every time. On-screen assisted calibration and verification save you time and ensure accuracy. With an easy-to-follow interface, complicated manuals are not needed to perform routine calibrations. Single-standard RapidCal<sup>™</sup> calibration offers a simplified solution for low level measurements.

### Simple Data Transfer

Data transfer with the optional USB + Power Module is simple, flexible, and doesn't require additional software. All data can be transferred to the module and easily downloaded to your computer with a USB connection, providing superior data integrity and availability. With two different module options, you can customize connectivity and power to meet your unique needs.

### Accurate for Rapidly Settling Samples

The Hach 2100Q Portable Turbidimeter incorporates an innovative Rapidly Settling Turbidity™ mode to provide accurate, repeatable measurements for difficult to measure, rapidly settling samples. An exclusive algorithm that

calculates turbidity based on a series of automatic readings eliminates redundant measurements and estimating.

### **Convenient Data Logging**

Up to 500 measurements are automatically stored in the instrument for easy access and backup. Stored information includes: date and time, operator ID, reading mode, sample ID, sample number, unlts, calibration time, calibration status, error messages and the result.

### **Optical System for Precision in the Fleld**

The two-detector optical system compensates for color in the sample, light fluctuation, and stray light, enabling analysts to achieve laboratory-grade performance on a wide range of samples, even under difficult site conditions.

### **Two Models for Specific Requirements**

- 2100Q Turbidimeter—Compliant with USEPA Method 180.1 design criteria.
- 2100Q is Turbidimeter—Compliant with ISO 7027 design criteria.

DW = drinking water WW = wastewater municipal PW = pure water / power IW = industrial water E = environmental C = collections FB = food and beverage



W P



### Specifications\*

Measurement Method Ratio turbidimetric determination using a primary nephelometric light scatter signal (90°) to the transmitted light scatter signal.

Regulatory 2100Q: Meets EPA Method 180.1 2100Q is: Meets ISO 7027

Light Source 2100Q: Tungsten filament lamp 2100Q is: Light-emitting diode (LED) @ 860 nm

Range 0 to 1000 NTU (FNU)

Accuracy ±2% of reading plus stray light from 0 to 1000 NTU

Repeatability  $\pm 1\%$  of reading, or 0.01 NTU (FNU), whichever is greater

Resolution 0.01 NTU on lowest range

Stray Light <0.02 NTU (FNU)

Signal Averaging Selectable on/off

Detector Silicon photovoltaic

Reading Modes (user selectable) Normal (Push to Read) Signal Averaging Rapidly Settling Turbidity

Data Logger 500 records

Power Requirement 110-230 Vac, 50/60 Hz (with Power or USB+Power Module) 4 AA alkaline batteries Rechargeable NiMH (for use with USB+Power Module) Operating Conditions Temperature: 0 to 50°C (32 to 122°F) Relative Humidity: 0 to 90% @ 30°C, 0 to 80% @ 40°C, 0 to 70% @ 50°C, noncondensing

Storage Conditions -40 to 60°C (-40 to 140°F), instrument only

Languages English, French, German, Italian, Spanish, Portuguese (BR), Portuguese (PT), Bulgarian, Chinese, Czech, Danish, Dutch, Finnish, Greek, Hungarian, Japanese, Korean, Polish, Romanian, Russian, Slovenian, Swedish, Turkish

Interface Optional USB

Instrument Enclosure Rating IP67 (closed lid, battery compartment excluded)

Protection Class Power Supply: Class II

Certification CE certified

Sample Required 15 mL (0.3 oz.)

Sample Cells 60 x 25 mm (2.36 x 1 in.) borosilicate glass with screw cap

*Dimensions* 22.9 x 10.7 x 7.7 cm (9.0 x4.2 x 3.0 in.)

*Weight* 527 g (1.16 lb) without batteries 618 g (1.36 lb) with four AA alkaline batteries

Warranty 1 year

\*Specifications subject to change without notice.

Appendix D

Stream:	Collector	ors:				
Location:	Date:	Date:				
RM:	Time:					
Lat/Long:						
Number of Rocks:	Total Area Scraped:	cm <sup>2</sup>				
		Diameter to Area Conversion				
Diameter of individual scrape	Area of individual scrape	Diameter (cm) Area (cm2)				
1	1	1.6 2.011				
2	2	1.7 2.27				
3	3	1.8 2.545				
4	4	1.9 2.835				
5	5	2.0 3.142				
6	6	2.1 3.464				
7	7	2.2 3.801				
8	8	2.3 4.155				
9	9					
10	10	Total Sample Volumeml				
11	11	Filter 1 LABLynx ID				
12	12	Volml				
13	13					
14	14	Filter 2 LABLynx ID				
15	15	Volml				
16	16					
17	17	Filter 3 LABLynx ID				
18	18	Volml				
19	19					
20	20					
21	21	Water Column Chlorophyll Sample				
22	22	Filter 1 LABLynx ID				
23	23	Volml				
24	24					
25	25	Filter 2 LABLynx ID				
	Total:	Volml				
		Filter 3 LABLynx ID				
		Volml				

### NEORSD Chlorophyll a Sampling Field Sheet

Flow:	None	Low	Normal	Elevated	High
<b>Turbidity:</b> *Explain	Clear	Low	Moderate*	High*	
Sky:	Overcast	Cloudy	Partly Cloudy	Mostly Clear	Clear
Canopy:	Open	Mostly Open	Partly Closed	Closed	
Riparian	None	Narrow L R	Moderate L R	Wide L R	

Downstream Channel Direction	Record two most predominate substrates with an X, and check all present.
$300^{\circ}$ $270^{\circ}$ W $240^{\circ}$ S $180^{\circ}$ $150^{\circ}$ $150^{\circ}$	RiffleRunReachBoulder/Slabs
Clinometer Left Bank° Right Bank°	Substrate Origin LimestoneTillsRip-rap SandstoneShaleWetlands LacustrineHardpanCoal Fines
Left Bank° Right Bank°	Silt HeavyModerateNormalNone
Left Bank° Right Bank°	Embeddedness ExtensiveModerateNormalNone
Stream Widths mmm Notes:	

Length of Reach: \_\_\_\_\_m

Stream Drawing

Appendix E

## **COMMONWEALTH OF PENNSYLVANIA**

# DEPARTMENT OF ENVIRONMENTAL PROTECTION

BUREAU OF LABORATORIES

LABORATORY ACCREDITATION PROGRAM

**Certifies** That

68-03670

Northeast Ohio Regional Sewer District Analytical Services 4747 East 49th Street, Cuyahoga Heights, OH 44125

DEPARTMENT OF ENVIRONMENTAL PROTECTION

pennsylvania

National Environmental Laboratory Accreditation Program Standard dealing with Environmental Laboratories Accreditation The act of June 29, 2002 (P.L. 596, No. 90) Having duly met the requirement of (27 Pa. C.S. §§4104-4113) and the

is hereby approved as an

### Accredited Laboratory

As more fully described in the attached Scope of Accreditation

Expiration Date: 11/30/2015 Certificate Number: 008

Laboratory Accreditation Program Aaren S. Alger, Chief Bureau of Laboratories

laun alger

Shall not be used to imply endorsement by the Commonwealth of Pennsylvania Not valid unless accompanied by a valid Scope of Accreditation To be conspicuously displayed at the Laboratory Certificate not transferable Surrender upon revocation Continued accreditation status depends on successful ongoing participation in the program

Customers are urged to verify the laboratory's current accreditation status

PA DEP is a NELAP recognized accreditation body





 Attached to Certificate of Accreditation 008-001 expiration date November 30, 2015. This listing of accredited analytes should be used only when associated with a valid certificate of accreditation.

 DEP Laboratory ID: 68-03670
 EPA Lab Code: OH00300
 TNI Code: (216) 641-6000

Northeast Ohio Regional Sewer District Analytical Services

4747 East 49th Street

Cuyahoga Heights, OH 44125

### Matrix: Non-Potable Water

Method	Revision	Analyte	Accreditation Type	Primary	Effective Date
ASTM D4839-03		Total organic carbon (TOC)	NELAP	PA	11/25/2014
EPA 1000.0		Pimephales promelas	NELAP	PA	1/8/2009
EPA 1002.0		Ceriodaphnia dubia	NELAP	PA	1/8/2009
EPA 160,4		Residue, volatile	NELAP	PA	10/22/2008
EPA 1600		Enterococci	NELAP	PA	11/22/2010
EPA 1603		E. coli (Enumeration)	NELAP	PA	11/29/2007
EPA 1631	E	Mercury	NELAP	PA	3/31/2008
EPA 180.1		Turbidity	NELAP	PA	12/31/2007
EPA 200.7	4.4	Aluminum	NELAP	PA	11/29/2007
EPA 200.7	4.4	Antimony	NELAP	PA	11/29/2007
EPA 200.7	4.4	Arsenic	NELAP	PA	11/29/2007
EPA 200.7	4.4	Barium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Beryllium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Cadmium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Calcium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Chromium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Cobalt	NELAP	PA	11/29/2007
EPA 200.7	4.4	Copper	NELAP	PA	12/31/2007
EPA 200.7	4.4	lron	NELAP	PA	11/29/2007
EPA 200.7	4.4	Lead	NELAP	PA	11/29/2007
EPA 200.7	4.4	Magnesium	NELAP	PA	11/17/2010
EPA 200.7	4.4	Manganese	NELAP	PA	11/29/2007
EPA 200.7	4.4	Molybdenum	NELAP	ΡA	11/29/2007
EPA 200.7	4.4	Nickel	NELAP	PA	11/29/2007
EPA 200.7	4.4	Potassium	NELAP	PA	12/31/2007
EPA 200.7	4.4	Selenium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Silver	NELAP	PA	11/29/2007
EPA 200.7	4.4	Sodium	NELAP	PA	12/31/2007
EPA 200.7	4.4	Thallium	NELAP	PA	4/15/2014
EPA 200.7	4.4	Tin	NELAP	PA	11/29/2007
EPA 200.7	4.4	Titanium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Vanadium	NELAP	PA	11/29/2007
EPA 200.7	4.4	Zinc	NELAP	PA	12/31/2007
EPA 245.1	3.0	Mercury	NELAP	PA	11/29/2007
EPA 300.0	2.1	Bromide	NELAP	PA	11/22/2010
EPA 300.0	2.1	Chloride	NELAP	PA	11/22/2010
EPA 300.0	2.1	Fluoride	NELAP	PA	11/22/2010
EPA 300.0	2.1	Nitrate as N	NELAP	PA	11/22/2010
EPA 300.0	2.1	Orthophosphate as P	NELAP	PA	11/22/2010
EPA 300.0	2.1	Sulfate	NELAP	PA	11/22/2010

Gener alger

The Pennsylvania Department of Environmental Protection Laboratory Accreditation Program is a NELAP recognized Accreditation Body. Customers are urged to verify the laboratory's current accreditation standing.

Page 1 of 4

www.dep.state.pa.us





Attached to Certificate of Accreditation 008-001 expiration date November 30, 2015. This listing of accredited analytes

should be used only when associated with a valid certificate of accreditation. DEP Laboratory ID: 68-03670 EPA Lab Code: OH00300 TNI Code: (216) 641-6000

### Matrix: Non-Potable Water

Method	Revision	Analyte	Accreditation Type	Primary	Effective Date
EPA 3005	А	Preconcentration under acid	NELAP	PA	11/29/2007
EPA 3010	A	Hot plate acid digestion (HNO3 + HCl)	NELAP	PA	11/29/2007
EPA 3015		Microwave-assisted acid digestion	NELAP	PA	11/29/2007
EPA 310.2		Alkalinity as CaCO3	NELAP	PA	9/20/2012
EPA 350 1		Ammonia as N	NELAP	PA	11/29/2007
EPA 351.2		Kjeldahl nitrogen, total (TKN)	NELAP	PA	11/17/2010
EPA 353.2		Nitrate as N	NELAP	PA	11/29/2007
EPA 353.2		Total nitrate-nitrite	NELAP	PA	11/17/2010
EPA 365 1		Phosphorus, total	NELAP	PA	10/22/2008
EPA 410,4		Chemical oxygen demand (COD)	NELAP	PA	11/29/2007
EPA 420.4		Total phenolics	NELAP	PA	11/17/2010
EPA 445		Chlorophyll A	NELAP	PA	11/22/2010
EPA 6010		Aluminum	NELAP	PA	11/29/2007
EPA 6010		Antimony	NELAP	PA	11/29/2007
EPA 6010		Arsenic	NELAP	PA	11/29/2007
EPA 6010		Barium	NELAP	PA	11/29/2007
EPA 6010		Beryllium	NELAP	PA	11/29/2007
EPA 6010		Cadmium	NELAP	PA	11/29/2007
EPA 6010		Calcium	NELAP	PA	11/29/2007
EPA 6010		Chromium	NELAP	PA	11/29/2007
EPA 6010		Cobalt	NELAP	PA	11/29/2007
EPA 6010		Copper	NELAP	PA	12/31/2007
EPA 6010		lron	NELAP	PA	11/29/2007
EPA 6010		Lead	NELAP	PA	11/29/2007
EPA 6010		Magnesium	NELAP	PA	11/29/2007
EPA 6010		Manganese	NELAP	PA	11/29/2007
EPA 6010		Molybdenum	NELAP	PA	11/29/2007
EPA 6010		Nickel	NELAP	PA	11/29/2007
EPA 6010		Potassium	NELAP	PA	12/31/2007
EPA 6010		Selenium	NELAP	PA	11/29/2007
EPA 6010		Silver	NELAP	PA	11/29/2007
EPA 6010		Sodium	NELAP	PA	12/31/2007
EPA 6010		Thallium	NELAP	PA	4/15/2014
EPA 6010		Tin	NELAP	PA	11/29/2007
EPA 6010		Titanium	NELAP	PA	11/29/2007
EPA 6010		Vanadium	NELAP	PA	11/29/2007
EPA 6010		Zinc	NELAP	PA	12/31/2007
EPA 7470		Mercury	NEL AP	PA	11/29/2007
Enternlert		Enterococci (Enumeration)	NELAP	PA	11/22/2010
014 1677		A vailable cvanide	NELAP	PA	11/29/2007
SM 2540 B		Residue total	NELAP	PA	11/29/2007
SM 2540 C		Residue, filterable (TDS)	NELAP	PA	11/29/2007
SM 2540 D		Residue, nonfilterable (TSS)	NELAP	PA	11/29/2007
SM 2540 F		Residue settleable	NELAP	PA	11/20/2001
SM 2550 P		Temperature deg	NELAP	PA	10/22/2007
SM 3500.Cr B	20-22	Chromium VI	NELAP	PA	11/20/2000
SM 4500-CN-C/F	20-22	Total evanide	NFLAP	PA	11/29/2001
AND A THIRD AND A		a named as welling that a share of the second	17 4-1-1 34	10	1 4/ 6 3/ 600

aun alger

The Pennsylvania Department of Environmental Protection Laboratory Accreditation Program is a NELAP recognized Accreditation Body. Customers are urged to verify the laboratory's current accreditation standing.

Page 2 of 4

www.dep.state.pa.us





Attached to Certificate of Accreditation 008-001 expiration date November 30, 2015. This listing of accredited analytes should be used only when associated with a valid certificate of accreditation.

DEP Laboratory ID: 68-03670 EPA I

EPA Lab Code: OH00300 TNI Code:

(216) 641-6000

### Matrix: Non-Potable Water

Method	Revision	Analyte	Accreditation Type	Primary	Effective Date
SM 4500-CN- G		Amenable cyanide	NELAP	PA	11/29/2007
SM 4500-C1 E		Total residual chlorine	NELAP	PA	11/29/2007
SM 4500-C1- C		Chloride	NELAP	PA	11/19/2012
SM 4500-H+ B		pH	NELAP	PA	11/29/2007
SM 4500-NO2- B		Nitrite as N	NELAP	PA	11/29/2007
SM 4500-Norg B		Kjeldahl nitrogen, total (TKN)	NELAP	PA	10/22/2008
SM 4500-P B	÷.	Preliminary treatment of phosphate samples	NELAP	PA	11/13/2013
SM 4500-P E		Orthophosphate as P	NELAP	PA	11/13/2013
SM 5210 B		Biochemical oxygen demand (BOD)	NELAP	PA	11/29/2007
SM 5210 B		Carbonaceous BOD (CBOD)	NELAP	PA	11/29/2007
SM 9222 D		Fecal coliform (Enumeration)	NELAP	PA	11/29/2007
SM 9223 Colilert MPN or QT		E. coli (Enumeration)	NELAP	PA	11/29/2007
SM 9223 Colilert MPN or QT		Total coliform (Enumeration)	NELAP	PA	11/22/2010

### Matrix: Solid and Chemical Materials

Method	Revision	Analyte	Accreditation Type	Primary	Effective Date
EPA 245.1	3.0	Mercury	NELAP	PA	11/22/2010
EPA 3051		Microwave digestion of solids (HNO3 only)	NELAP	PA	11/17/2010
EPA 6010		Aluminum	NELAP	PA	11/22/2010
EPA 6010		Antimony	NELAP	PA	11/13/2013
EPA 6010		Arsenic	NELAP	PA	11/22/2010
EPA 6010		Barium	NELAP	PA	11/22/2010
EPA 6010		Beryllium	NELAP	PA	11/22/2010
EPA 6010		Cadmium	NELAP	PA	11/22/2010
EPA 6010		Calcium	NELAP	PA	11/22/2010
EPA 6010		Chromium	NELAP	PA	11/22/2010
EPA 6010		Cobalt	NELAP	PA	11/22/2010
EPA 6010		Copper	NELAP	PA	11/22/2010
EPA 6010		Iron	NELAP	PA	11/22/2010
EPA 6010		Lead	NELAP	PA	11/22/2010
EPA 6010		Magnesium	NELAP	PA	11/22/2010
EPA 6010		Manganese	NELAP	PA	11/22/2010
EPA 6010	В	Metals by ICP/AES	NELAP	PA	1/22/2013
EPA 6010		Molybdenum	NELAP	PA	11/22/2010
EPA 6010		Nickel	NELAP	PA	11/22/2010
EPA 6010		Potassium	NELAP	PA	11/22/2010
EPA 6010		Selenium	NELAP	PA	11/22/2010
EPA 6010		Silver	NELAP	PA	11/22/2010
EPA 6010		Sodium	NELAP	PA	11/22/2010
EPA 6010		Thallium	NELAP	PA	11/22/2010
EPA 6010		Tin	NELAP	PA	4/15/2013
EPA 6010		Titanium	NELAP	PA	11/22/2010
EPA 6010		Vanadium	NELAP	PA	11/22/2010
EPA 6010		Zinc	NELAP	PA	11/22/2010

aven alger

The Pennsylvania Department of Environmental Protection Laboratory Accreditation Program is a NELAP recognized Accreditation Body. Customers are urged to verify the laboratory's current accreditation standing.

Page 3 of 4

www.dep.state.pa.us





TNI Code:

Attached to Certificate of Accreditation 008-001 expiration date November 30, 2015. This listing of accredited analytes should be used only when associated with a valid certificate of accreditation.

DEP Laboratory ID: 68-03670

EPA Lab Code: OH00300

(216) 641-6000

Gen alger

The Pennsylvania Department of Environmental Protection Laboratory Accreditation Program is a NELAP recognized Accreditation Body. Customers are urged to verify the laboratory's current accreditation standing.

Page 4 of 4

www.dep.state.pa.us

Appendix F



February 4, 2015

Mr. John Rhoades Supervisor of Environmental Assessment Northeast Ohio Regional Sewer District 4747 East 49<sup>th</sup> Street Cuyahoga Heights, Ohio 44125

Dear Mr. Rhoades:

This letter is to acknowledge that I am responsible for assisting the Northeast Ohio Regional Sewer District's Water Quality and Industrial Surveillance Division in conducting stream habitat assessments using the Qualitative Habitat Evaluation Index for the 2015 Big Creek, Cuyahoga River, Euclid Creek, Hemlock Creek, Mill Creek, and West Creek Environmental Monitoring Project Study Plans.

It is understood that an Ohio Environmental Protection Agency Level 3 Qualified Data Collector Certification for Stream Habitat Assessment is required to perform these tasks and that I am responsible for maintaining my Level 3 QDC Certification during the term of these Study Plans.

In addition, I have not been convicted nor pleaded guilty to a Violation of Section 2911.21 of the Revised Code (criminal trespass) or a substantially similar municipal ordinance within the previous five years.

Sincerely

Jonathan Brauer Stormwater Inspector Northeast Ohio Regional Sewer District 4747 East 49<sup>th</sup> Street Cuyahoga Heights, Ohio 44125



April 14, 2015

Mr. John Rhoades Manager of Water Quality and Industrial Surveillance Northeast Ohio Regional Sewer District 4747 East 49<sup>th</sup> Street Cuyahoga Heights, Ohio 44125

Re: 2015 Benthic Services PO 15001329

Dear Mr. Rhoades:

This letter is to acknowledge that I am responsible for the identification of benthic macroinvertebrates for the following Northeast Ohio Regional Sewer District Study Plans:

- 2015 Big Creek Environmental Monitoring
- 2015 Cuyahoga River Environmental Monitoring
- 2015 Euclid Creek Environmental Monitoring
- 2015 Hemlock Creek Environmental Monitoring
- 2015 Mill Creek Environmental Monitoring
- 2015 West Creek Environmental Monitoring

It is understood that an Ohio Environmental Protection Agency Level 3 Qualified Data Collector Certification for Benthic Macroinvertebrate, with the specialty of identification, is required to perform these tasks and that I am responsible for maintaining my Level 3 QDC Certification during the term of these Study Plans.

In addition, I have not been convicted nor pleaded guilty to a Violation of section 2911.21 of the Revised Code (criminal trespass) or a substantially similar municipal ordinance within the previous five years.

Sincerely,

albert W. Kember I

Bert Remley Senior Taxonomist Bremley@thirdrockconsultants.com

THIRD ROCK CONSULTANTS, LLC, 2526 REGENCY ROAD, SUITE 180, LEXINGTON, KY 40503 PHONE 859.977.2000 • FAX 859.977.2001 • WWW.THIRDROCKCONSULTANTS.COM Appendix G

Will send Wildlife Permit upon Receipt Appendix H

### References

Chlorophyll a Sampling and Field Filtering Standard Operating Procedure (SOP-EA001-00)

- EPA New England- Region 1. (2005). Standard operating procedure for calibration and field measurement procedures for the YSI Model 6-Series Sondes and Data Logger (Including: temperature, pH, specific conductance, turbidity, dissolved oxygen, chlorophyll, rhodamine WT, ORP, and barometric pressure) (7<sup>th</sup> Revision). North Chelmsford, MA: The Office of Environmental Measurement and Evaluation, Ecosystem Assessment- Ecology Monitoring Team.
- Ohio Environmental Protection Agency. (1987a). Biological criteria for the protection of aquatic life: Volume II. Users manual for biological field assessment of Ohio surface waters (Updated January 1988; September 1989; November 2006; August 2008). Columbus, OH: Division of Water Quality Monitoring and Assessment.
- Ohio Environmental Protection Agency. (1987b). Biological criteria for the protection of aquatic life: Volume III. Standardized biological field sampling and laboratory methods for assessing fish and macroinvertebrate communities (Updated September 1989; March 2001; November 2006; and August 2008). Columbus, OH: Division of Water Quality Monitoring and Assessment.
- Ohio Environmental Protection Agency. (1997). Draft. Biological Criteria for the Protection of Aquatic Life: Volume IV: Fish and Macroinvertebrate Indicies for Ohio's Lake Erie Nearshore Waters, Harbors, and Lacustuaries. Columbus, OH: Division of Surface Water, Ecological Assessment Unit.
- Ohio Environmental Protection Agency. (2006). Methods for assessing habitat in flowing waters: using the Qualitative Habitat Evaluation Index (QHEI). (Ohio EPA Technical Bulletin EAS/2006-06-1). Columbus, OH: Division of Surface Water; Division of Ecological Assessment Section.
- Ohio Environmental Protection Agency. (2010). Draft. *Methods of Assessing Habitat in Lake Erie Shoreline Waters Using the Qualitative Habitat Evaluation Index (QHEI) Approach (Version 2.1)*. Columbus, OH: Division of Surface Water.
- Ohio Environmental Protection Agency. (2011). State of Ohio Water Quality Standards Ohio Administrative Code Chapter 3745-1 (Revision: March 16, 2011; Effective June 16, 2011). Columbus, OH: Division of Surface Water; Standards and Technical Support Section.
- Ohio Environmental Protection Agency. (2012a). *Field Evaluation Manual for Ohio's Primary Headwater Habitat Stream.* Columbus, OH: Division of Surface Water; Division of Ecological Assessment Section.

Ohio Environmental Protection Agency. (2012b). Ohio 2012 Integrated Water Quality Monitoring and Assessment Report. Columbus, Ohio: Division of Surface Water.

Ohio Environmental Protection Agency. (2013). Surface Water Field Sampling Manual for water chemistry, bacteria, and flows. Columbus, OH: Division of Surface Water.